

-1-

5

ENHANCED TRIPLE-HELIX AND DOUBLE-HELIX FORMATION  
WITH OLIGOMERS CONTAINING MODIFIED PYRIMIDINES

10 This is a continuation-in-part of United States  
Patent Application attorney docket number 24610-20035.21,  
filed October 23, 1992, which is a continuation-in-part  
of United States Patent Application Serial No.  
07/935,444, filed 25 August 1992, which is a  
continuation-in-part of United States Patent Application  
15 Serial No. 799,824, filed 26 November 1991, and each  
incorporated herein by reference.

Technical Field

20 The invention relates generally to novel  
nucleomonomer and oligomer analogs, and to  
oligonucleotide-based therapy and diagnosis by binding of  
the oligonucleotide analogs to single or double-stranded  
nucleic acid target sequences. More specifically, the  
invention concerns oligomers containing certain 5-  
25 substituted pyrimidine base residues and intermediates in  
their synthesis.

Background Art

30 Sequence-specific binding of oligonucleotides  
both to single-stranded RNA and DNA and to duplex DNA has  
been demonstrated. The appropriate sequence recognition  
for binding to single-stranded targets is well known:  
the A-T and G-C pairing characteristic of duplex  
35

-2-

formation has been established as the basis for DNA replication and transcription.

More recently, oligonucleotides have been shown to bind in a sequence-specific manner to duplex DNA to form triplexes. Single-stranded nucleic acid, primarily RNA, is the target molecule for oligonucleotides that are used to inhibit gene expression by an "antisense" mechanism (Uhlmann, E., et al, Chem Reviews (1990) 90:543-584; van der Krol, A.R., et al, Biotechniques (1988) 6:958-976). Antisense oligonucleotides are postulated to exert an effect on target gene expression by hybridizing with a complementary RNA sequence. In this model, the hybrid RNA-oligonucleotide duplex interferes with one or more aspects of RNA metabolism including processing, translation and metabolic turnover. Chemically modified oligonucleotides have been used to enhance their nuclease stability.

Duplex DNA can be specifically recognized by oligomers based on a recognizable nucleomonomer sequence. Two major recognition motifs have been recognized. In an earlier description of a "CT" motif, protonated cytosine residues recognize G-C basepairs while thymine residues recognize A-T basepairs in the duplex. These recognition rules are outlined by Maher III, L.J., et al., Science (1989) 245:725-730; Moser, H.E., et al., Science (1987) 238:645-650. More recently, an additional motif, termed "GT" recognition, has been described (Beal, P.A., et al, Science (1992) 251:1360-1363; Cooney, M., et al., Science (1988) 241:456-459; Hogan, M.E., et al., EP Publication 375408). In the G-T motif, A-T pairs are recognized by adenine or thymine residues and G-C pairs by guanine residues.

35

-3-

In both of these binding motifs, the recognition sequence of the oligomer must align with the complementary sequence of the purine chain of the duplex; thus, recognition, for example, of an A-T pair by a thymine, depends on the location of the adenyl residues along the purine chain of the duplex. An exception to the foregoing is the recent report by Griffin, L.C., et al., Science (1989) 245:967-971, that limited numbers of guanine residues can be provided within pyrimidine-rich oligomers and specifically recognize thymine-adenine base pairs; this permits the inclusion of at least a limited number of pyrimidine residues in the homopurine target.

The two motifs exhibit opposite binding orientations with regard to homopurine target chains in the duplex. In the CT motif, the targeting oligonucleotide is oriented parallel to the target purine-rich sequence; in the GT motif, the oligonucleotide is oriented antiparallel (Beal, P.A., et al., Science (1990) 251:1360-1363).

The efficiency of binding by C residues in CT motif oligomers is reduced as the pH of hybridization is increased. The protonated tautomer of C (C<sup>+</sup>) is the binding competent species in Hoogsteen binding, but is present at only low levels at physiological pH. This is consonant with the pK<sub>a</sub> of cytosine which is 4.25. Base analogs such as 5-methylcytosine, pK<sub>a</sub> 4.35, (Lee, J.S. et al., Nucleic Acids Res (1984) 12:6603-6614), 8-oxo-N<sup>6</sup>-methyladenine (Krawczyk, S.H. et al, Proc Natl Acad Sci (1992) 89:3761-3764; International Application No. PCT/US91/08811), pseudoisocytidine (Ono, A., et al, J Org Chem (1992) 57:3225-3230; International Application No. PCT/US90/03275) or carbocyclic cytidine (Froehler, B.C., et al, J Am Chem Soc (1992) 114:8320-8322; U.S. Patent

-4-

Application Serial No. 07/864,873 incorporated herein by reference) have been utilized to obtain binding of CT motif oligomers over an extended pH range.

Sequence-specific targeting of both single-stranded and duplex target sequences has applications in diagnosis, analysis, and therapy. Under some circumstances wherein such binding is to be effected, it is advantageous to stabilize the resulting duplex or triplex over long time periods.

Covalent crosslinking of the oligomer to the target provides one approach to prolong stabilization. Sequence-specific recognition of single-stranded DNA accompanied by covalent crosslinking has been reported by several groups. For example, Vlassov, V.V., et al., Nucleic Acids Res (1986) 14:4065-4076, describe covalent bonding of a single-stranded DNA fragment with alkylating derivatives of nucleomonomers complementary to target sequences. A report of similar work by the same group is that by Knorre, D.G., et al., Biochimie (1985) 67:785-789. Iverson and Dervan also showed sequence-specific cleavage of single-stranded DNA mediated by incorporation of a modified nucleomonomer which was capable of activating cleavage (J Am Chem Soc (1987) 109:1241-1243). Meyer, R.B., et al., J Am Chem Soc (1989) 111:8517-8519, effect covalent crosslinking to a target nucleomonomer using an alkylating agent complementary to the single-stranded target nucleomonomer sequence. Photoactivated crosslinking to single-stranded oligonucleotides mediated by psoralen was disclosed by Lee, B.L., et al., Biochemistry (1988) 27:3197-3203. Use of crosslinking in triple-helix forming probes was also disclosed by Horne, et al., J Am Chem Soc (1990) 112:2435-2437.

35



-5-

Use of N<sup>4</sup>,N<sup>4</sup>-ethan cytosine as an alkylating agent to crosslink to single-stranded and double-stranded oligomers has also been described (Webb and Matteucci, J Am Chem Soc (1986) 108:2764-2765; Nucleic Acids Res (1986) 14:7661-7674; Shaw, J.P., et al, J Am Chem Soc (1991) 113:7765-7766). These papers also describe the synthesis of oligonucleotides containing the derivatized cytosine. Matteucci and Webb, in a later article in Tetrahedron Letters (1987) 28:2469-2472, describe the synthesis of oligomers containing N<sup>6</sup>,N<sup>6</sup>-ethanoadenine and the crosslinking properties of this residue in the context of an oligonucleotide binding to a single-stranded DNA.

In a recent paper, Praseuth, D., et al., Proc Natl Acad Sci (USA) (1988) 85:1349-1353, described sequence-specific binding of an octathymidylate conjugated to a photoactivatable crosslinking agent to both single-stranded and double-stranded DNA.

In addition, Vlassov, V.V. et al., Gene (1988) 313-322 and Fedorova, O.S. et al., FEBS (1988) 228:273-276, describe targeting duplex DNA with an alkylating agent linked through a 5'-phosphate of an oligonucleotide.

In effecting binding to obtain a triplex, to provide for instances wherein purine residues are concentrated on one chain of the target and then on the opposite chain, oligomers of inverted polarity can be provided. By "inverted polarity" is meant that the oligomer contains tandem sequences which have opposite polarity, i.e., one having polarity 5'→3' followed by another with polarity 3'→5', or vice versa. This implies that these sequences are joined by linkages which can be thought of as effectively a 3'-3' internucleoside

-6-

junction (however the linkage is accomplished), or  
effectively a 5'-5' internucleoside junction. Such  
oligomers have been suggested as by-products of reactions  
to obtain cyclic oligonucleotides by Capobionco, M.L., et  
5 al., Nucleic Acids Res (1990) 18:2661-2669. Compositions  
of "parallel-stranded DNA" designed to form hairpins  
secured with AT linkages using either a 3'-3' inversion  
or a 5'-5' inversion have been synthesized by van de  
Sande, J.H., et al., Science (1988) 241:551-557. In  
10 addition, triple helix formation using oligomers which  
contain 3'-3' linkages have been described (Horne, D.A.,  
and Dervan, P.B., J Am Chem Soc (1990) 112:2435-2437;  
Froehler, B.C., et al, Biochemistry (1992) 31:1603-1609).

The use of triple helix (or triplex) complexes  
15 as a means for inhibition of the expression of target  
gene expression has been previously adduced  
(International Application No. PCT/US89/05769). Triple  
helix structures have been shown to interfere with target  
gene expression (International Application No.  
20 PCT/US91/09321; Young, S.L. et al, Proc Natl Acad Sci  
(1991) 88:10023-10026), demonstrating the feasibility of  
this approach.

European Patent Application No. 92103712.3,  
Rahim, S.G., et al (Antiviral Chem Chemother (1992)  
25 3:293-297), and International Application No.  
PCT/SE91/00653 describe pyrimidine nucleomonomer  
characterized by the presence of an unsaturated group in  
the 5-position. Propynyl and ethynyl groups are included  
among the derivatives at the 5-position that are  
30 described in the applications.

Synthesis of nucleomonomers having unsaturated  
alkyl groups at the 5-position of uracil has been  
described (DeClercq, E., et al, J Med Chem (1983) 26:661-  
35

-7-

666; Goodchild, J., et al, J Med Chem (1983) 26:1252-1257). Oligom rs containing 5-pr pynyl modified pyrimidines have been described (Froehler, B.C., et al, Tetrahedron Letters (1992) 33:5307-5310).

5                   Conversion of 5-propynyl-2'-deoxyuridine, 5-butynyl-2'-deoxyuridine and related compounds to the 5'-triphosphate followed by incorporation of the monomer into oligomers by E. coli polymerase has been described (Valko, K., et al, J Liquid Chromatog (1989) 12:2103-  
10 2116; Valko, K. et al, J Chromatog (1990) 506:35-44). These studies were conducted as a structure to activity analysis of nucleotide analogs having a series of substitutions at the 5- position of uracil. The activity of the nucleotide analogs as substrates for E. coli  
15 polymerase was examined and correlated with characteristics such as the hydrophobicity of the monomer. No information was presented regarding the properties of oligomers containing the analogs.

                  European patent application 0492570 published  
20 July 1, 1992 describes a method for detecting a target polynucleotide using a single-stranded polynucleotide probe in which an intercalating molecule is attached by a linker which comprises at least 3 carbon atoms and a double bond at the alpha position relative to the base.

25                   PCT patent publication WO 92/02258 describes nuclease resistant, pyrimidine modified oligomers including a substituent group at the 5 or 6 positions, including phenyl. 5-Phenyl-2'-deoxyuridine has been subsequently incorporated into oligomers and shown to  
30 decrease the binding affinity of oligomers containing this modification for both single stranded and double stranded target sequences.

35

-8-

DNA synthesis via amidite and hydrogen phosphonate chemistries has been described (U.S. Patent Nos. 4,725,677; 4,415,732; 4,458,066; 4,959,463).

Oligomers having enhanced affinity for complementary target nucleic acid sequences would have improved properties for diagnostic applications, therapeutic applications and research reagents. Thus, a need exists for oligomers with enhanced binding affinity for complementary sequences. Oligomers of the present invention have improved binding affinity for double stranded and/or single stranded target sequences.

Brief Description of Figures

1. Dimer synthons containing bases of the invention.
2. Dimer synthons containing bases of the invention and containing 5 and 6 membered riboacetal type linkages.
3. Dimer synthons containing bases of the invention and containing 6 and 7 membered riboacetal type linkages.
4. Synthesis of monomers containing 2'-O-allyl modifications.
5. Synthesis of o-xylene linked switchback dimers (synthetic method #1).
6. Synthesis of monomers containing 2'-S-alkyl modifications.
7. Synthesis of dimer linked by a 3'-thioformacetal linkage (method #2).
8. Synthesis of trimer linked by a 3'-thioformacetal linkage (method #2).
9. Synthesis of dimer linked by a riboacetal linkage (method #3).

-9-

10. Coupling groups for oligomer synthesis via amidite or triester chemistry.

11. Synthesis of dimer linked by a formacetal linkage (method #2).

5 12. Oligomer synthesis by (1) hydrogen-phosphonate, (2) amidite chemistry and (3) methyl phosphonate derivatives (method #1).

13. Synthesis of a monomer for an oligomer containing amide linkages (method #4).

10 14. Synthesis of the 5-((1-ethynyl)-2-pyrimidinyl)-2'-deoxyuridine nucleomonomer.

15 15. Synthesis of 5-(2-pyridinyl)-2'-deoxyuridine and 5-(2-pyridinyl)-2'-deoxycytidine nucleomonomers.

16 16. Synthesis of 5-(2-thienyl)-2'-deoxyuridine derivative.

17. Oligomers containing amide substitute linkages; repeating nucleomonomer units and exemplary amide-linked oligomer structures.

20 18. Dimers synthons containing bases of the invention and having exemplary 2',5' linkages; thioformacetal and carbamate linkages.

#### Structural Formulas

25 Structural formulas that are described herein are designated as a numeral in parentheses ((1), (2), etc.) and chemical compounds are designated as an underlined numeral (1, 2, etc.).

#### 30 Disclosure of the Invention


The invention provides an oligomer comprising at least two and preferably a multiplicity of

35


-10-

nucleomonomers wherein at least one nucleomonomer  
comprises a base of formula (1) or (2)

5



10



wherein each X is independently O or S;

15

R<sup>2</sup> is a group comprising at least one pi bond  
connected to a carbon atom attached to the base; and

Pr is (H), or a protecting group,

with the proviso that when at least one of said  
nucleomonomers of said oligomer comprises deoxyuridine 5-  
substituted by vinyl, 1-butenyl, 1-pentenyl, 1-hexenyl,  
1-heptenyl, 1-octenyl, 1-propynyl, 1-butyryl, 1-hexynyl,  
1-heptynyl, or 1-octynyl, then the remainder of the  
nucleomonomers comprising said oligomer are not solely  
comprised of phosphodiester linked 2'-deoxyadenosine, 2'-  
deoxyguanosine, 2'-deoxycytidine, thymidine or a  
combination thereof.

25

### Definitions

The following definitions are brief synopses of  
terms that are more fully defined hereinbelow.

30

Nucleomonomer. As used herein, the term  
"nucleomonomer" means a moiety comprising (1) a base  
covalently linked to (2) a second moiety. Nucleomonomers  
include nucleosides and nucleotides. Nucleomonomers can  
be linked to form oligomers that bind to target r

35

-11-

complementary base sequences in nucleic acids in a sequence specific manner.

5 A "second moiety" as used herein includes those species which contain modifications of the sugar moiety, for example, wherein one or more of the hydroxyl groups are replaced with a halogen, a heteroatom, an aliphatic groups, or are functionalized as ethers, amines, and the like. The pentose moiety can be replaced by a hexose or an alternate structure such as a cyclopentane ring, a 6-  
10 member morpholino ring and the like. Nucleosides as defined herein are also intended to include a base linked to an amino acid and/or an amino acid analog having a free carboxyl group and/or a free amino group and/or protected forms thereof

15 Base. "Base" as used herein includes those moieties which contain not only the known purine and pyrimidine heterocycles and the invention pyrimidines, but also heterocycle analogs and tautomers thereof. Purines include adenine, guanine and xanthine and  
20 exemplary purine analogs include 8-oxo-N<sup>6</sup>-methyladenine and 7-deazaxanthine. Pyrimidines include uracil and cytosine and their analogs such as 5-methylcytosine, 5-methyluracil and 4,4-ethanocytosine. Invention bases are pyrimidines derivatized at the 5- position. The  
25 derivatives are 1-alkenyl-, 1-alkynyl-, heteroaromatic- and 1-alkynyl-heteroaromatic modifications. "1-Alkenyl" means an olefinically-unsaturated (double bond containing) acyclic group. "1-Alkynyl" means an acetylenically-unsaturated (triple bond containing)  
30 acyclic group. "Heteroaromatic" means a compound having at least one heterocyclic ring, 5 or 6 ring atoms, having physical and chemical properties resembling compounds such as an aromatic group. "Heteroaromatic" also means  
35 systems having one or more rings, including bicyclic moieties such as benzimidazole, benzotriazole,

-12-

benzoxazole, and indole. A base also includes heterocycles such as 2-aminopyridine and triazines. 1-Alkynyl-heteroaromatic means 1-ethynyl-heteroaryl wherein heteroaryl is as defined above.

5            Nucleoside. As used herein, "nucleoside" means a base covalently attached to a sugar or sugar analog and which may contain a phosphite or phosphine. The term nucleoside includes ribonucleosides, deoxyribonucleosides, or any other nucleoside which is an  
10 N-glycoside or C-glycoside of a base. The stereochemistry of the sugar carbons can be other than that of D-ribose.

Nucleosides include those species which contain modifications of the sugar moiety, for example, wherein  
15 one or more of the hydroxyl groups are replaced with a halogen, a heteroatom, an aliphatic groups, or are functionalized as ethers, amines, thiols, and the like. The pentose moiety can be replaced by a hexose or an alternate structure such as a cyclopentane ring, a 6-  
20 member morpholino ring and the like. Nucleosides as defined herein are also intended to include a base linked to an amino acid and/or an amino acid analog having a free carboxyl group and/or a free amino group and/or protected forms thereof.

25            Nucleotide. As used herein, "nucleotide" means nucleoside having a phosphate group or phosphate analog.

Sugar Modification. As used herein, "sugar modification" means any pentose or hexose moiety other than 2'-deoxyribose. Modified sugars include D-ribose,  
30 2'-O-alkyl, 2'-amino, 2'-halo functionalized pentoses, hexoses and the like.. Sugars having a stereochemistry other than that of a D-ribose are also included.

35            Linkage. As used herein, "linkage" means a phosphodiester moiety (-O-P(O)(O)-O-) that covalently couples adjacent nucleomonomers.



-13-

Substitute Linkages. As used herein, "substitute linkage" means any analog of the native phosphodiester group that covalently couples adjacent nucleomonomers. Substitute linkages include

5 phosphodiester analogs, e.g. such as phosphorothioate and methylphosphonate, and nonphosphorus containing linkages, e.g. such as acetals and amides.

Switchback. As used herein, "switchback" means an oligomer having at least one region of inverted polarity. Switchback oligomers are able to bind to  
10 opposite strands of a duplex to form a triplex on both strands of the duplex. The linker joining the regions of inverted polarity is a substitute linkage.

Oligomers. Oligomers are defined herein as two  
15 or more nucleomonomers covalently coupled to each other by a linkage or substitute linkage moiety. Thus, an oligomer can have as few as two nucleomonomers (a dimer). Oligomers can be binding competent and, thus, can base pair with cognate single-stranded or double-stranded  
20 nucleic acid sequences. Oligomers (e.g. dimers - hexamers) are also useful as synthons for longer oligomers as described herein. Oligomers can also contain abasic sites and pseudonucleosides.

Blocking Groups. As used herein, "blocking group" refers to a substituent other than H that is  
25 conventionally attached to oligomers or nucleomonomers, either as a protecting group, a coupling group for synthesis,  $PO_3^{2-}$ , or other conventional conjugate such as a solid support. As used herein, "blocking group" is not  
30 intended to be construed solely as a protecting group, according to slang terminology, but also includes, for example, coupling groups such as a hydrogen phosphonate or a phosphoramidite.

35 Protecting group. "Protecting group" as used herein means any group capable of preventing the O-atom

-14-

or N-atom to which it is attached from participating in a reaction or bonding. Such protecting groups for O- and N-atoms in nucleomonomers are described and methods for their introduction are conventionally known in the art.  
5 Protecting groups also prevent reactions and bonding at carboxylic acids, thiols and the like.

Coupling group. "Coupling group" as used herein means any group suitable for generating a linkage or substitute linkage between nucleomonomers such as a  
10 hydrogen phosphonate and a phosphoramidite.

Conjugate. "Conjugate" as used herein means any group attached to the oligomer at a terminal end or within the oligomer itself. Conjugates include solid supports, such as silica gel, controlled pore glass and  
15 polystyrene; labels, such as fluorescent, chemiluminescent, radioactive, enzymatic moieties and reporter groups; oligomer transport agents, such as polycations, serum proteins and glycoproteins and polymers and the like.

20 Pi bond. "Pi bond" as used herein means an unsaturated covalent bond such as a double or triple bond. Both atoms can be carbon or one can be carbon and the other nitrogen, for example, phenyl, propynyl, cyano and the like.

25 Synthon. "Synthon" as used herein means a structural unit within a molecule that can be formed and/or assembled by known or conceivable synthetic operations.

30 Transfection. "Transfection" as used herein refers to any method that is suitable for enhanced delivery of oligomers into cells.

35 Subject. "Subject" as used herein means an animal, including a mammal, particularly a human.

-15-

Description of the Invention

Oligomers including either or both of the modified bases (1) or (2) show enhanced binding capacities in the formation of duplexes or triplexes with single-stranded RNA or DNA or duplex target sequences, respectively.

When the certain 5-substituted pyrimidines noted above are present, the additional nucleomonomer modifications can vary widely as discussed hereinafter. Preferably, the additional modification is at least one substitute linkage or a sugar modification such as a 2'-substituted deoxyribose.

The substitution of a base (1) or (2) of the invention, such as in 5-(1-alkenyl)-, 5-(1-alkynyl)-, 5-heteroaromatic- or 1-alkynyl-heteroaromatic substituted bases for thymine or cytosine in oligomers which target DNA duplexes provides binding competent oligomers with enhanced binding affinity. Substitution for thymine base residues by the 5-substituted uracil or thiouracil base residues of the invention or substitution for cytosine or 2-thiocytosine base residues by the 5-substituted cytosine base residues of the invention enhance the ability of the resulting oligomer to bind single-stranded DNA or RNA targets. In addition, some of the 5-substituted pyrimidine base residues significantly enhance triple helix formation with double stranded DNA.

For some  $R^2$ , substitution of 5- $R^2$  substituted U (5- $R^2$ -U) for T in oligomers results in enhanced ability to form triplexes and duplexes as compared with the oligomers containing thymine. 5- $R^2$ -U in these oligomers, in triplex formation recognizes adenine residues in adenine-thymine base pairs when hybridized in the parallel CT triplex motif. Oligomers having 8-oxo- $N^6$ -methyladenine (a cytosine analog for triplex binding) and 5- $R^2$ -U also bind in the CT motif. Oligomers having 5- $R^2$ -U

-16-

and guanine are suitable for triplex binding to duplex  
sequences via the GT motif (5-R<sup>2</sup>-U recognizes adenine).  
Some oligomers containing substitution of 5-R<sup>2</sup>  
substituted C (5-R<sup>2</sup>-C) in place of C bind duplex DNA, but  
5 not as well as control oligomers containing  
5-methylcytosine at corresponding positions. The reduced  
efficiency of triplex formation is believed to result  
primarily from the reduced pK<sub>a</sub> of the substituted base.  
In the 5-propynyl-substituted nucleomonomer corresponding  
10 to the nucleomonomer containing 5-methylcytosine, the pK<sub>a</sub>  
is only 3.30.

The oligomers of the invention are thus capable  
of forming triplexes with various target sequences such  
as those found in oncogenes or viruses by binding in the  
15 major groove of a target DNA duplex under physiological  
pH conditions.

However, alteration of the heterocycle pK<sub>a</sub> as  
described above for the 5-R<sup>2</sup>-C does not significantly  
affect binding to single-stranded target nucleic acid.  
20 In addition to binding efficiently to double-stranded  
target sequences, oligomers of the invention containing  
5-R<sup>2</sup> substituted U in place of T and/or 5-R<sup>2</sup> substituted C  
in place of C were also found to bind single-stranded RNA  
efficiently. Oligomers containing either 5-R<sup>2</sup>-C or 5-R<sup>2</sup>-U  
25 formed duplex structures with complementary single-  
stranded RNA that had increased thermal stability (T<sub>m</sub>)  
compared to the duplex formed by a control oligomer as  
described below.

Accordingly, in one aspect, the invention is  
30 directed to an oligomer comprising at least two and  
preferably, a multiplicity, of nucleomonomers wherein at  
least one said nucleomonomer comprises a base of formula  
35 (1) or (2) above.

-17-

Preferably, each X is O, i.e. formula (1) is uracil and formula (2) is cytosine. Other suitable pyrimidines include 4-thiouracil, 2-thiouracil, 2,4-dithiouracil and 2-thiocytosine.

5 In one embodiment of the invention R<sup>2</sup> is cyano, C<sub>2-12</sub> 1-alkenyl or 1-alkynyl or is a C<sub>2-12</sub> heteroaromatic group containing 5-6 ring atoms in which one to three of the ring atoms is N, S or O. Preferably, R<sup>2</sup> is a C<sub>2-4</sub> 1-alkenyl or 1-alkynyl or a C<sub>2-4</sub> heteroaromatic group  
10 containing 5-6 ring atoms in which one ring atom is N and optionally a second ring atom is N, S or O.

By "1-alkenyl" is meant an olefinically-unsaturated acyclic group, for example, vinyl, 1-propenyl, 1-butenyl optionally substituted by halogen or  
15 an alkynyl group.

By "1-alkynyl" is meant an acetylenically-unsaturated acyclic group, such as ethynyl, 1-propynyl, 1-butylnyl, 1-pentylnyl, 1,3-pentadiynyl, and the like optionally substituted by an aryl or heteroaryl group,  
20 such as phenylethynyl, pyridine-ethynyl, pyrimidine-ethynyl, triazine-ethynyl, thiophene-ethynyl, thiazole-ethynyl and imidazole-ethynyl.

By "heteroaromatic" is meant a compound having at least one heterocyclic ring having physical and  
25 chemical properties resembling compounds such as an aromatic group of from 5 to 6 ring atoms and 2 to 12 carbon atoms in which one to three ring atoms is N, S or O, for example, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 2-thiazoyl, triazinyl, 2-imidazolyl, 2-oxazolyl, 2-pyridinyl (o-pyridinyl), 3-pyridinyl (m-pyridinyl), 4-pyridinyl (p-pyridinyl), 2-thienyl, 2-furanyl, 2-pyrrolyl optionally substituted preferably on  
30 a ring C by oxygen, alkyl of 1-4 carbon atoms or halogen  
35

-18-

or on a ring N by alkyl of 1-4 carbon atoms. Preferred substituents on the heteroaryl group are methyl, ethyl, trifluoromethyl, and bromo.

Preferred oligomers contain one or more 5-R<sup>2</sup>-U  
5 or 5-R<sup>2</sup>-C bases.

In another embodiment of the invention, the oligomer comprises at least one base of formula (1) or (2) wherein each X is independently O or S; and R<sup>2</sup> is selected from the group consisting of phenylethynyl, 2-,  
10 3-, and 4-pyridine-ethynyl, 2-, 4- and 5-pyrimidine-ethynyl, triazine-ethynyl, 2-, 4- and 5-pyrimidinyl, 2-, 4- and 5-thiazolyl, 1-methyl-2-imidazolyl, 2- and 4-imidazolyl, 2-, 4- and 5-oxazolyl, 3-pyridinyl, 4-pyridinyl, 2-pyridinyl, 2- and 3-furanyl-ethynyl, 2- and  
15 3-thienyl-ethynyl, 2- and 4-imidazolyl-ethynyl, 2-, 4- and 5-thiazolyl-ethynyl, 2-, 4- and 5-oxazolyl-ethynyl, 2- and 3-pyrrolyl-ethynyl, 2- and 3-thienyl, 2- and 3-furanyl, 2- and 3-pyrrolyl, propenyl (-CH=CH-CH<sub>3</sub>), vinyl and -C≡C-Z where Z is hydrogen (H) or C<sub>1-10</sub> alkyl,  
20 haloalkyl (C<sub>1-10</sub> with 1 to 6 halogen atoms or heteroalkyl (C<sub>1-10</sub> with 1 to 3 heteroatoms selected from the group consisting of O, N and S); and

Pr is (H), or a protecting group.

Examples of -C≡C-Z include 1-propynyl (-C≡C-CH<sub>3</sub>), 3-buten-1-ynyl (-C≡C-CH=CH<sub>2</sub>), 3-methyl-1-butynyl (-C≡C-CH(CH<sub>3</sub>)<sub>2</sub>), 3,3-dimethyl-1-butynyl (-C≡C-C(CH<sub>3</sub>)<sub>3</sub>), 1-butynyl (-C≡C-CH<sub>2</sub>-CH<sub>3</sub>), 1,3-pentadiynyl (-C≡C-C≡C-CH<sub>3</sub>) and ethynyl.

Preferred halogens are selected from the group  
30 consisting of fluorine, chlorine and bromine.  
Substitutions including bromovinyl can be included in the oligomers.


35

-19-

Aspects of the invention include the use of nucleomonomers, two linked nucleomonomers (dimers), three linked nucleomonomers (trimers), four linked nucleomonomers (tetramers), five linked nucleomonomers (pentamers) or six linked nucleomonomers (hexamers) as intermediates in the synthesis of the longer oligomers of the invention. These oligomers are valuable synthons of the invention that are useful in the synthesis of longer oligomers.

In other aspects, the invention is directed to duplexes or triplexes obtained by binding the foregoing oligomers to single-stranded or duplex nucleic acid targets.

Other useful intermediates in the synthesis of the oligomers of the invention include an o-xyloso dimer having the structural formula (5),



(5)

wherein each Y is independently an oligomer or R<sup>1</sup>; and each B is independently a base provided that at least one B is a base of formula (1) or (2) wherein R<sup>2</sup> is as defined herein.

Also included are intermediates of the formula (6) shown in Figure 1, wherein X is selected from the group consisting of O and S; and each Y, B, and R<sup>3</sup> is

-20-

independently selected and has the meaning defined herein.

5 The oligomers of the invention are also suitable for binding to DNA duplex target sequences via either CT or GT triple helix binding motif.

10 The novel oligomers of the present invention are useful in antisense therapies wherein an oligomer hybridizes with a selected complementary RNA sequence or triple helix therapies wherein an oligomer hybridizes with a selected complementary DNA sequence.

15 The invention is also directed to an oligomer of the invention comprising a positive modification of least one base of formula (1) or (2) of the invention and a negative modification, each with respect to the binding affinity of the oligomer to a complementary nucleic acid sequence. The positive modification counteracts the effect of the negative modification to a degree that is more than additive with respect to the binding affinity, thus a synergistic effect is observed.

20 An aspect of the invention is the inclusion of the invention bases in oligomers that are resistant to nuclease degradation relative to an oligodeoxynucleotide having no modifications. Nuclease resistant oligomers of the invention are advantageously used under conditions where nucleases are present. For certain applications, such as modulation of gene expression by via an antisense mechanism, nuclease stability by oligomers of the invention is an important functional aspect of the oligomer.

30 Other aspects of the invention are directed to pharmaceutical compositions, reagents and kits comprising the oligomers of the invention, to methods of treating conditions, such as cancers and viruses or the like.

35



-21-

Such conditions are associated with or characterized by particular nucleic acids such as DNA duplexes or single-stranded RNA or DNA.

5 An additional aspect of the invention includes methods of detecting the presence, absence or amount of a particular single-stranded DNA or RNA or a particular target duplex in a biological (or other) sample using the oligomers of the invention, to detect selected nucleic acid sequences. Such sequences can be associated with  
10 the presence of neoplastic growth, viruses or disease conditions. Reagents and kits containing oligomers of the invention represent an aspect of the invention that permit facile use of the oligomers as reagents useful for (1) modulating gene expression in cells in vitro  
15 including cells grown in tissue culture, and (2) detecting and/or quantitating target sequences.

It has been found that some of the oligomers of the invention have enhanced binding properties with respect to complementary single-stranded and double-  
20 stranded nucleic acid sequences as compared to unmodified oligomers not having the 5-substitution of the invention. Triple helix structures can be formed at physiological pH levels of 7.0 and higher, where unmodified control oligomers were less efficient. Improved duplex formation  
25 is also noted.

A feature of the invention is that the oligomers of the invention can be comprised of a variety of different sequences and thereby used to target a variety of different single-stranded or double-stranded  
30 target sequences.

An advantage of the present invention is that the oligomers of the invention are capable of forming triplexes under physiological pH conditions.  
35

-22-

Another feature of oligomers containing 5-R<sup>2</sup> substituted uracil or cytosin base (1) or (2) of the invention compared to oligomers containing thymine or cytosine is that the lipophilic group (R<sup>2</sup>) can enhance cell permeation or uptake. The nucleomonomers containing these bases are more lipophilic than uridine, cytidine or thymidine based on retention times on reverse phase HPLC.

10 Additional Nucleomonomer Modifications.

Oligomers that are comprised of nucleomonomers can also contain modifications in addition to the 5-modified pyrimidines of the invention. A non-limiting exemplary list of such additional modifications includes oligomers where (i) one or more nucleomonomer residues are modified at the 2' position, (ii) one or more covalent crosslinking moieties are incorporated, (iii) inverted polarity linkers are incorporated, (iv) substitute linkages are included, (v) other base analogs, such as 8-oxo-N<sup>6</sup>-methyladenine, are included and (vi) conjugates such as intercalating agents or polylysine that respectively enhance binding affinity to target nucleic acid sequences or that enhance association of the oligomer with cells are included.

25 The ability of the 5-substitution of the bases (1) and (2) of the invention to enhance affinity of the oligomer for single-stranded and duplex targets (positive modification) permits further modifications to the oligomer in which they are contained. These further modifications may or may not diminish affinity, but also confer other useful properties such as stability to nuclease cleavage, ability to permeate cell membranes, and the like. Any decrease in binding affinity resulting

-23-

from the further modifications (negative modification) is acceptable because of the enhanced affinity conferred by the 5-substituted bases (1) and (2). Thus, particularly preferred oligomers of the invention can contain  
5 substitute linkages and/or modified sugars, as well as the 5-substituted pyrimidine bases (1) and (2) of the invention.

The oligomers can also contain additional modifications in the nucleomonomers that contain these 5-  
10 modified pyrimidines or in other nucleomonomers that comprise the oligomer.

Also included are oligomers containing one or more substitute linkages such as sulfide or sulfone linkages (Benner, S.A., International Publication No. WO  
15 89/12060), sulfamate linkages (International Publication No. WO 91/15500), carbamate linkages in morpholino-linked oligomers (Stirchak, E.P. et al Nucleic Acids Res (1989) 17:6129-6141) and related linkages in morpholino oligomers of the formula (7) shown in Figure 1  
20 wherein  $X^2$  is CO, CS or  $SO_2$ ;  $X^3$  is O, S, NH,  $NCH_3$ ,  $CH_2$ ,  $CF_2$  or CHF; each Y is independently an oligomer or  $R^1$  and each B is independently chosen and has the previously defined meaning, provided that at least one B is a base of formula (1) or (2).

25 Riboacetal and related linkages, amide linkages and 2',5' linkages are described in commonly owned pending U.S. application serial numbers 07/806,710, filed December 12, 1991, 07/899,736, filed Can 28, 1992, 07/894,397 and filed 07/892,902, filed each cited  
30 reference is incorporated herein by reference.

Exemplary dimers containing riboacetal and related linkages of formulae (8-15) are shown in Figures  
35 2 and 3 wherein for each structure,

-24-

R<sup>1</sup> and B are independently chosen and have the meanings defined above;

R<sup>3</sup> has the meaning as defined above;

5 X<sup>3</sup> is independently selected from the group consisting of O, S, NH, NCH<sub>3</sub>, CH<sub>2</sub>, CF<sub>2</sub>, and CFH;

X<sup>4</sup> is independently selected from the group consisting of O, S, SO, SO<sub>2</sub>, CH<sub>2</sub>, CO, CF<sub>2</sub>, CS, NH and NR<sup>4</sup> wherein R<sup>4</sup> is lower alkyl (C<sub>1-4</sub>; methyl, ethyl, propyl, isopropyl, butyl or isobutyl);

10 X<sup>5</sup> is selected from the group consisting of O, CO, S, CH<sub>2</sub>, CS, NH and NR<sup>4</sup>;

X<sup>6</sup> is selected from the group consisting of CH, N, CF, CCl, and CR<sup>5</sup> wherein R<sup>5</sup> is methyl or lower alkyl (C<sub>2-4</sub>) fluoromethyl, difluoromethyl, trifluoromethyl or lower fluoroalkyl (C<sub>2-4</sub>, F<sub>1-3</sub>);

15 X<sup>7</sup> is selected from the group consisting of O, S, CH<sub>2</sub>, CO, CF<sub>2</sub>, and CS,

provided that at least one B is of the formula (1) or (2) as defined above; and

20 further provided that no adjacent X<sup>4</sup>, X<sup>5</sup> or X<sup>7</sup> are O (i.e., -O-O-, a peroxide).

Compounds of the 5-member ring series are preferred embodiments for oligomers containing one or more riboacetal linkages (formula (8)), where X<sup>4</sup> is O and X<sup>5</sup>, X<sup>7</sup> are CH<sub>2</sub> and X<sup>6</sup> is CH.

Also included are oligomers containing nucleomonomer residues linked via amide bonds. Exemplary linkages have been described (Nielsen, P.E., et al, Science (1991) 254:1497-1500; commonly owned copending U.S. Application Serial Nos. 07/889,736, filed Jan 28, 1992, and 07/894,397, filed June 5, 1992, both incorporated herein by reference).

35

-25-

Oligomers

As used herein "oligomer" includes oligonucleotides, oligonucleosides, polydeoxyribonucleotides (containing 2'-deoxy-D-ribose or modified forms thereof), i.e., DNA, polyribonucleotides (containing D-ribose or modified forms thereof), i.e., RNA, and any other type of polynucleotide which is an N-glycoside or C-glycoside of a purine or pyrimidine base, or modified purine or pyrimidine base. Oligomer as used herein is also intended to include compounds where adjacent nucleomonomers are linked via amide linkages as previously described (Nielsen, P.E., et al, Science (1991) 254:1497-1500). The enhanced competence of binding by oligomers containing the bases of the present invention is believed to be primarily a function of the base alone. Because of this, elements ordinarily found in oligomers, such as the furanose ring and/or the phosphodiester linkage can be replaced with any suitable functionally equivalent element. "Oligomer" is thus intended to include any structure that serves as a scaffold or support for the bases wherein the scaffold permits binding to target nucleic acids in a sequence-dependent manner. Oligomers that are currently known can be defined into four groups that can be characterized as having (i) phosphodiester and phosphodiester analog (phosphorothioate, methylphosphonate, etc) linkages, (ii) substitute linkages that contain a non-phosphorous isostere (formacetal, riboacetal, carbamate, etc), (iii) morpholino residues, carbocyclic residues or other furanose sugars, such as arabinose, or a hexose in place of ribose or deoxyribose and (iv) nucleomonomers linked via amide bonds or acyclic nucleomonomers linked via any suitable substitute linkage.

35

-26-

The oligomers of the invention can be formed using invention and conventional nucleomonomers and synthesized using standard solid phase (or solution phase) oligomer synthesis techniques, which are now commercially available. In general, the invention oligomers can be synthesized by a method comprising the steps of: synthesizing a nucleomonomer or oligomer synthon having a protecting group and a base and a coupling group capable of coupling to a nucleomonomer or oligomer; coupling the nucleomonomer or oligomer synthon to an acceptor nucleomonomer or an acceptor oligomer; removing the protecting group; and repeating the cycle as needed until the desired oligomer is synthesized.

The oligomers of the present invention can be of any length including those of greater than 40, 50 or 100 nucleomonomers. In general, preferred oligomers contain 2-30 nucleomonomers. Lengths of greater than or equal to about 8 to 20 nucleomonomers are useful for therapeutic or diagnostic applications. Short oligomers containing 2, 3, 4 or 5 nucleomonomers are specifically included in the present invention and are useful as synthons.

Oligomers having a randomized sequence and containing about 6 or 7 nucleomonomers are useful for primers that are used in cloning or amplification protocols that use random sequence primers, provided that the oligomer contains residues that can serve as a primer for polymerases or reverse transcriptases.

Oligomers can contain conventional phosphodiester linkages or can contain substitute linkages such as phosphoramidate linkages. These substitute linkages include, but are not limited to, embodiments wherein a moiety of the formula  $-O-P(O)(S)-O-$

-27-

("ph sphorothioate"),  $-O-P(S)(S)-O-$   
("ph sphorodithioat "),  $-O-P(O)(NR'_2)-X-$ ,  $-O-P(O)(R')-O-$ ,  
 $-O-P(S)(R')-O-$  ("thionoalkylphosphonate"),  $-P(O)(OR^6)-X-$ ,  
 $-O-C(O)-X-$ , or  $-O-C(O)(NR'_2)-X-$ , wherein  $R'$  is H (or a  
5 salt) or alkyl (1-12C) and  $R^6$  is alkyl (1-9C) and the  
linkage is joined to adjacent nucleomonomers through an -  
O- or -S- bonded to a carbon of the nucleomonomer.  
Phosphorothioate and phosphodiester linkages are shown in  
Figure 12. Particularly, preferred substitute linkages  
10 for use in the oligomers of the present invention include  
phosphodiester, phosphorothioate, methylphosphonate and  
thionomethylphosphonate linkages. Phosphorothioate and  
methylphosphonate linkages confer added stability to the  
oligomer in physiological environments. While not all  
15 such linkages in the same oligomer need be identical,  
particularly preferred oligomers of the invention contain  
uniformly phosphorothioate linkages or uniformly  
methylphosphonate linkages.

## 20 Pharmaceutically Acceptable Salts

Any pharmaceutically acceptable salt can be  
used and such salt forming materials are well known in  
the art.

Pharmaceutically acceptable salts are  
25 preferably metal or ammonium salts of said oligomers of  
the invention and include alkali or alkaline earth metal  
salts, e.g., the sodium, potassium, magnesium or calcium  
salt; or advantageously easily crystallizing ammonium  
salts derived from ammonia or organic amines, such as  
30 mono-, di- or tri-lower (alkyl, cycloalkyl or  
hydroxyalkyl)-amides, lower alkylenediamines or lower  
(hydroxyalkyl or arylalkyl)-alkylammonium bases, e.g.  
methylamine, diethylamine, triethylamine,  
35

-28-

dicyclohexylamine, triethan lamine, ethylen diamine, tris-(hydroxymethyl)-aminomethane or benzyl-trimethylammonium hydroxide. The oligomers of the invention form acid addition salts, which are preferably  
5 such of therapeutically acceptable inorganic or organic acids, such as strong mineral acids, for example hydrohalic, e.g., hydrochloric or hydrobromic acid; sulfuric, phosphoric; aliphatic or aromatic carboxylic or sulfonic acids, e.g., formic, acetic, propionic,  
10 succinic, glycollic, lactic, malic, tartaric, gluconic, citric, ascorbic, maleic, fumaric, hydroxymaleic, pyruvic, phenylacetic, benzoic, 4-aminobenzoic, anthranilic, 4-hydroxybenzoic, salicylic, 4-aminosalicylic, methanesulfonic, ethanesulfonic,  
15 hydroxyethanesulfonic, benzenesulfonic, sulfanilic or cyclohexylsulfamic acid and the like.

#### Blocking Groups

As used herein, "blocking group" refers to a  
20 substituent other than H that is conventionally coupled to oligomers or nucleomonomers, either as a protecting group, a coupling group for synthesis,  $\text{PO}_3^{2-}$ , or other conventional conjugate such as a solid support, label, antibody, monoclonal antibody or fragment thereof and the  
25 like. As used herein, "blocking group" is not intended to be construed solely as a protecting group, according to slang terminology, but is meant also to include, for example, coupling groups such as a H-phosphonate or a phosphoramidite.

30 By "protecting group" is meant is any group capable of protecting the O-atom or N-atom to which it is attached from participating in a reaction or bonding. Such protecting groups for N-atoms on a base moiety in a  
35



-29-

nucleomonomer and their introduction are conventionally known in the art. Non-limiting examples of suitable protecting groups include diisobutylformamidine, benzoyl and the like. Suitable "protecting groups" for O-atoms  
5 are, for example, DMT, MMT, or FMOC.

Suitable coupling groups are, for example, H-phosphonate, a methylphosphonamidite, or a phosphoramidite. Phosphoramidites that can be used include  $\beta$ -cyanoethylphosphoramidites (preferred).  
10 Methylphosphonamidites, alkylphosphonamidites (including ethylphosphonamidites and propylphosphonamidites) can also be used. Exemplary phosphoramidites are shown in Figures 10-1 and 10-2.

Suitable protecting groups are DMT (dimethoxy trityl), MMT (monomethoxytrityl) or FMOC at the 5' terminus and/or hydrogen phosphonate, methyl phosphoramidite, methyl phosphonamidite,  $\beta$ -cyanoethylphosphoramidite at the 3'-terminus.

## 20 Protecting Groups

Protecting groups such as diisobutylformamidine, benzoyl, isobutyryl, FMOC, dialkylformamidine, dialkylacetamidine or other groups known in the art can be used to protect the exocyclic  
25 nitrogen of the cytosine heterocycle. Alternatively, cytidine precursors can be directly incorporated into oligomers without a protecting group at the exocyclic nitrogen using described methods (Gryaznov, S.M. et al, J Amer Chem Soc (1991) 113:5876-5877; Gryaznov, S.M., et  
30 al, Nucl Acids Res (1992) 20:1879-1882; Kung, P.-P., et al, Tetrahedron Letters (1992) 40:5869-5872). Synthesis of oligomers having bases (1) or (2) containing an R<sup>2</sup> as ethynyl heteroaryl substituents is preferably  
35

-30-

accomplished using 9-fluorenylmethoxycarbonyl (FMOC) for protection of the 5'-hydroxyl position as described (Lehman, C., et al, Nucl Acids Res (1989) 17:2379-2390).

Preferred protecting groups are DMT (dimethoxy trityl), MMT (monomethoxytrityl) or FMOC at the 5' terminus and/or hydrogen phosphonate, methyl phosphoramidite, methyl phosphonamidite,  $\beta$ -cyanoethylphosphoramidite at the 3'- terminus. However, it is intended that the position of the blocking groups can be reversed as needed (e.g., a phosphoramidite at the 5'- position and DMT at the 3'- position). In general, the nucleomonomers and oligomers of the invention can be derivatized to such "blocking groups" as indicated in the relevant formulas by methods known in the art.

#### Coupling Groups

Suitable coupling groups are, for example, H-phosphonate, a methylphosphonamidite, or a phosphoramidite. Phosphoramidites that can be used include  $\beta$ -cyanoethylphosphoramidites (preferred). Methylphosphonamidites, alkylphosphonamidites (including ethylphosphonamidites and propylphosphonamidites) can also be used. Exemplary phosphoramidites are shown in Figures 10-1 and 10-2. Suitable "coupling groups" at the 3', 2' or 5' position for oligomer synthesis via phosphoramidite triester chemistry, referred to herein as "amidite" chemistry, include N,N-diisopropylamino- $\beta$ -cyanoethoxyphosphine, N,N-diisopropylamino-methoxyphosphine, N,N-diethylamino- $\beta$ -cyanoethoxyphosphine, (N-morpholino)- $\beta$ -cyanoethoxyphosphine, and (N-morpholino)-methoxyphosphine (Moore, M.F. et al, J Org Chem (1985) 50:2019-2025; Uznanski, A.W., et al, Tet Lett (1987) 28:3401-3404;

-31-

Bjergarde, K., et al, Nucl Acids Res (1991) 21:5843-5850;  
Dahl, O. Sulfur Reports (1991) 11:167-192). Related  
coupling groups such as N,N-diisopropylamino-methyl-  
phosphine or N,N-diethylamino-methyl-phosphine can also  
5 be used to prepare methylphosphonates (Fig 10-4).  
Methylphosphonate oligomers can be conveniently  
synthesized using coupling groups such as N,N-  
diisopropylamino-methylphosphonamidite, and N,N-  
diethylamino-methylphosphonamidite. Synthesis of  
10 nucleomonomer amidites of the invention can be  
accomplished by conventional methods (for example,  
Gryaznov, S.M., et al, Nucl Acids Res (1992) 20:1879-  
1882; Vinayak, R., et al, Nucl Acids Res (1992) 20:1265-  
1269; Sinha, N.D., et al, Nucl Acids Res (1984) 12:4539-  
15 4557; and other references cited herein). Suitable  
coupling groups at the 3', 2' (or 5') position for  
oligomer synthesis via phosphate triester chemistry,  
referred to herein as "triester" chemistry, include 2-  
chlorophenyl phosphate, 4-chlorophenyl phosphate, 2,4-  
20 dichlorophenyl phosphate and 2,4,-dibromophenyl phosphate  
nucleotide diester derivatives or, for synthesis of  
phosphorothioate linkages, the thiono derivatives thereof  
(Marugg, J.E., et al, Nucl Acids Res (1984) 12:9095-9110;  
Kemal, O., et al, J Chem Soc Chem Commun (1983) 591-593;  
25 Kamer, P.C.J., et al, Tet Lett (1989) 30:6757-6760).  
Structures of these coupling groups are shown in Figure  
10 where X is O or S and Z' is H or a suitable  
benzotriazole.

Oligomers or the segments thereof are  
30 conventionally synthesized. The synthetic methods known  
in the art and described herein can be used to synthesize  
oligomers containing bases of the invention, as well as  
other bases known in the art, using appropriately  
35

-32-

protected nucleomonomers (see Figure 12). Methods for the synthesis of oligomers are found, for example, in Froehler, B., et al., Nucleic Acids Res (1986) 14:5399-5467; Nucleic Acids Res (1988) 16:4831-4839; Nucleosides and Nucleotides (1987) 6:287-291; Froehler, B., Tetrahedron Letters (1986) 27:5575-5578; Caruthers, M. H. in Oligodeoxynucleotides-Antisense Inhibitions of Gene Expression (1989), J. S. Cohen, editor, CRC Press, Boca Raton, p7-24; Reese, C.B. et al, Tetrahedron Letters (1985) 26:2245-2248. Synthesis of the methylphosphonate linked oligomers via methyl phosphoramidite chemistry has also been described (Agrawal, S. et al., Tetrahedron Letters (1987) 28:3539-3542; Klem, R. E., et al, International Publication Number WO 92/07864).

#### Conjugates

Also included are "conjugates" of oligomers. "Conjugates" of the oligomers include those conventionally recognized in the art. For instance, the oligomers can be covalently linked to various moieties such as, intercalators, and substances which interact specifically with the minor groove of the DNA double helix. Other chosen conjugate moieties, can be labels such as radioactive, fluorescent, enzyme, or moieties which facilitate cell association using cleavage linkers and the like. Suitable radiolabels include  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^3\text{H}$  and  $^{14}\text{C}$ ; and suitable fluorescent labels include fluorescein, resorufin, rhodamine, BODIPY (Molecular Probes) and texas red; suitable enzymes include alkaline phosphatase and horseradish peroxidase. Other compounds which can be used as covalently linked moieties include biotin, antibodies or antibody fragments, transferrin and

-33-

the HIV Tat protein can also conveniently be linked to the oligomers of the invention.

These additional moieties can be derivatized through any convenient linkage. For example, intercalators, such as acridine or psoralen can be linked to the oligomers of the invention through any available -OH or -SH, e.g., at the terminal 5'- position of the oligomer, the 2'- positions of RNA, or an OH, NH<sub>2</sub>, COOH or SH incorporated into the 5- position of pyrimidines. A derivatized form which contains, for example, -CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH or -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>SH in the 5- position is preferred. Conjugates including polylysine or lysine can be synthesized as described and can further enhance the binding affinity of an oligomer to its target nucleic acid sequence (Lemaitre, M. et al., Proc Natl Acad Sci (1987) 84:648-652; Lemaitre, M. et al., Nucleosides and Nucleotides (1987) 6:311-315).

A wide variety of substituents can be attached, including those bound through linkages or substitute linkages. The -OH moieties in the oligomers can be replaced by phosphate groups, protected by standard protecting groups, or coupling groups to prepare additional linkages to other nucleomonomers, or can be bound to the conjugated substituent. The 5'- terminal OH can be phosphorylated; the 2'-OH or OH substituents at the 3'- terminus can also be phosphorylated. The hydroxyls can also be derivatized to standard protecting groups.

Oligomers of the invention can be covalently derivatized to moieties that facilitate cell association using cleavable linkers. Linkers used for such conjugates can include disulfide linkages that are reduced after the oligomer-transport agent conjugate has

-34-

entered a cell. Appropriate molecular linkers include  
for example,  $-Y^1-X^8CH_2CHR^7-SS-CHR^7CH_2X^8-Y^1-$  wherein each  $Y^1$   
is independently alkylene ( $C_{1-6}$ ; including methylene,  
ethylene and propylene), or CO, each  $X^8$  is independently  
5 O, S(O)(O), S(O),  $NR^7$ ,  $CH_2$ ,  $C(R^7)_2$ , or CO;  $R^7$  wherein each  $R^7$   
is independently H, alkyl ( $C_{1-6}$ ; including methyl, ethyl  
and propyl), or aryl and which linkers have been  
previously described (WO 91/14696). Disulfide-containing  
linkers of this type have a controllable  $t_{1/2}$  in vivo,  
10 facilitating its use as a prodrug/transport component.  
Such linkers are stable under extracellular conditions  
relative to intracellular conditions due to the redox  
potential of the disulfide linkage.

Suitable conjugates also include solid supports  
15 for oligomer synthesis and to facilitate detection of  
nucleic acid sequences. Solid supports included, but are  
not limited to, silica gel, controlled pore glass,  
polystyrene, and magnetic glass beads.

20

#### Sugar Modifications.

Derivatives can be made by substitution on the  
sugars. Among the most preferred derivatives of the  
oligomers of the invention are the 2'-O-allyl  
25 derivatives. The presence of the 2'-O-allyl group  
appears to enhance permeation ability and stability to  
nuclease degradation, but does not appear to diminish the  
affinity of the oligomer for single chain or duplex  
targets.

30

Furthermore, as the  $\alpha$  anomer binds to duplex  
DNA or single-stranded RNA in a manner similar to that  
for the  $\beta$  anomers but with a reversed polarity, oligomers  
can contain nucleomonomers having this epimer or a domain  
35

-35-

th reof (Praseuth, D., et al., Proc Natl Acad Sci (USA)  
(1988) 85:1349-1353; Sun, J.S. et al, Proc Natl Acad Sci  
(1991) 88:6023-6027; Debart, F., et al, Nucl Acids Res  
(1992) 20:1193-1200).  $\alpha$ -Anomeric oligomers containing  
5 the 5-R<sup>2</sup> substituted pyrimidines described herein  
represent a class of modified oligomers included in the  
present invention.

#### Substitute Linkages

10 The oligomers of the invention can also contain  
one or more "substitute linkages" as is generally  
understood in the art. These "substitute linkages"  
include phosphorothioate, methylphosphonate,  
thionomethylphosphonate, phosphorodithioate, riboacetal,  
15 2',5' linkages, alkylphosphonates, morpholino carbamate,  
morpholino sulfamate, morpholino sulfamide,  
boranophosphate (-O-P(OCH<sub>3</sub>)(BH<sub>3</sub>)-O-), siloxane (-O-  
Si(X<sup>4</sup>)(X<sup>4</sup>)-O-; X<sup>4</sup> is alkyl or phenyl) and phosphoramidate  
(methoxyethylamine and the like), and are synthesized as  
20 described in the generally available literature and in  
references cited herein (Sood, A., et al, J Am Chem Soc  
(1990) 112:9000-9001; WO 91/08213; WO 90/15065; WO  
91/15500; Stirchak, E.P. et al Nucleic Acid Res (1989)  
17:6129-6141; U.S. Patent 5,034,506; U.S. Patent  
25 5,142,047; Hewitt, J.M. et al, Nucleosides and  
Nucleotides (1992) 11:1661-1666). Substitute linkages  
that can be used in the oligomers disclosed herein also  
include the sulfonamide (-O-SO<sub>2</sub>-NH-), sulfide  
(-CH<sub>2</sub>-S-CH<sub>2</sub>-), sulfonate (-O-SO<sub>2</sub>-CH<sub>2</sub>-), carbamate  
30 (-O-C(O)-NH-, -NH-C(O)-O-), dimethylhydrazino  
(-CH<sub>2</sub>-NCH<sub>3</sub>-NCH<sub>3</sub>-), sulfamate (-O-S(O)(O)-N-;  
-N-S(O)(O)-N-), 3'-thioformacetal (-S-CH<sub>2</sub>-O-),  
formacetal (-O-CH<sub>2</sub>-O-), 3'-amine (-NH-CH<sub>2</sub>-CH<sub>2</sub>-),  
35

-36-

N-methylhydroxylamine ( $-\text{CH}_2-\text{NCH}_3-\text{O}-$ )  
and 2',5' linkages (such as 2',5' carbamate  
(2'  $-\text{N}(\text{H})-\text{C}(\text{O})-\text{O}-5'$ ), 5',2' carbamate (2'  
 $-\text{O}-\text{C}(\text{O})-\text{N}(\text{H})-5'$ ), 5',2' methylcarbamate (2'  
5  $-\text{O}-\text{C}(\text{O})-\text{N}(\text{CH}_3)-5'$ ) and 5',2' thioformacetal (2'  
 $-\text{O}-\text{CH}_2-\text{S}-5'$ ). 2',5' linkages are disclosed in pending  
U.S. application serial No. 07/892,902, filed June 1,  
1992, incorporated herein by reference). Riboacetal  
linkages are disclosed and claimed in commonly owned  
10 pending U.S. patent application serial nos. 690,786,  
filed April 24, 1991, 763,130, filed September 20, 1991,  
and 806,710 filed December 12, 1991, incorporated herein  
by reference. Except where specifically indicated, the  
substitute linkages, such as a formacetal linkage,  $-\text{O}-$   
15  $\text{CH}_2-\text{O}-$ , are linked to either the 3' or 2' carbon of a  
nucleomonomer on the left side and to the 5' carbon of a  
nucleomonomer on the right side. A formacetal linked  
(3',5') dimer is shown in Figure 1, formula (6). Thus a  
formacetal linkage can be indicated as 3'  $-\text{O}-\text{CH}_2-\text{O}-5'$  or  
20 2'  $-\text{O}-\text{CH}_2-\text{O}-5'$ . The designations of a 3', 2' or 5'  
carbon can be modified accordingly when a structure other  
than ribose, deoxyribose or arabinose is linked to an  
adjacent nucleomonomer. Such structures include a  
hexose, morpholino ring, carbocyclic ring (e.g.  
25 cyclopentane) and the like.

The use of carbamate, carbonate, sulfide,  
sulfoxide, sulfone, N-methylhydroxylamine and  
dimethylhydrazino linkages in synthons or oligomers has  
been described (Vaseur, J-J. et al, J Amer Chem Soc  
30 (1992) 114:4006-4007; WO 89/12060; Musicki, B. et al, J  
Org Chem (1990) 55:4231-4233; Reynolds, R. C., et al J  
Org Chem (1992) 57:2983-2985; Mertes, M. P., et al, J Med  
Chem (1969) 12:154-157; Mungall, W. S., et al, J Org Chem  
35



-37-

(1977) 42:703-706; Stirchak, E. P., et al, J Org Chem  
(1987) 52:4202-4206; Wang, H., et al, Tet Lett (1991)  
50:7385-7388; International Application No. PCT

US91/03680). Substitute linkage(s) can be utilized in  
5 the oligomers for a number of purposes such as to further  
facilitate binding with complementary target nucleic acid  
sequences and/or to increase the stability of the  
oligomers toward nucleases.

By "positive modification" is meant use of any  
10 modification of formula (1) or (2) of the invention which  
results in increased binding affinity.

By "negative modification" is meant an  
additional nucleomonomer modification of an oligomer  
comprising a base of formula (1) or (2) which results in  
15 a decrease in binding affinity or use of a substitute  
linkage which may result in a decrease in binding  
affinity.

For example, a negative substitute linkage  
modification is at least one selected from  
20 phosphorothioate, methylphosphonate,  
thionomethylphosphonate, phosphoroamidate and triester  
for a phosphodiester linkage.

#### Nucleosides

25 The term "nucleoside" will include  
ribonucleosides, deoxyribonucleosides, or to any other  
nucleoside which is an N-glycoside or C-glycoside of a  
purine or pyrimidine base, or modified purine or  
pyrimidine base. The stereochemistry of the sugar  
30 carbons can be other than that of D-ribose in one or more  
residues. The pentose moiety can be replaced by a hexose  
and incorporated into oligomers as described (Augustyns,  
K., et al Nucl Acids Res (1992) 18:4711-4716). Also  
35

-38-

included are analogs where the ribose or deoxyribose moiety is replaced by an alternate structure such as a hexose or such as the 6-member morpholino ring described in U.S. patent number 5,034,506. Nucleosides as defined  
5 herein also includes a purine or pyrimidine base linked to an amino acid or amino acid analog having a free carboxyl group and a free amino group or protected forms thereof. Exemplary nucleosides have been described (Nielsen, P.E. *ibid*; commonly owned copending U.S.  
10 Application Serial Nos. 07/889,736, filed Can 28, 1992, and 07/894,397, filed June 5, 1992, both applications incorporated herein by reference in their entirety).

"Nucleosides" also include those moieties which contain modifications of the sugar, for example, wherein  
15 one or more of the hydroxyl groups are replaced with halogen, aliphatic groups, or functionalized as ethers, amines, and the like. Such structures include a hexose, morpholino ring, carbocyclic ring (e.g. cyclopentane) and the like.

20

#### Base

"Base" as used herein includes those moieties which contain not only the known purine and pyrimidine bases and the invention bases, but also heterocyclic  
25 bases which have been modified and tautomers thereof. Such modifications include alkylated purines or pyrimidines, acylated purines or pyrimidines, or other heterocycles. Such "analogous purines" and "analogous pyrimidines" or purine or pyrimidine analogs are those  
30 generally known in the art, some of which are used as chemotherapeutic agents. An exemplary, but not exhaustive, list includes N<sup>4</sup>,N<sup>4</sup>-ethanocytosine, 7-deazaxanthosine, 7-deazaguanosine, 8-oxo-N<sup>6</sup>-  
35

-39-

methyladenine, 4-acetylcytosine, 5-  
(carb xyhydroxylmethyl) uracil, 5-fluorouracil,  
5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil,  
5-carboxymethylaminomethyl uracil, dihydrouracil,  
5 inosine, N<sup>6</sup>-isopentenyl-adenine, 1-methyladenine,  
1-methylpseudouracil, 1-methylguanine, 1-methylinosine,  
2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-  
methylcytosine, 5-methylcytosine, N<sup>6</sup>-methyladenine, 7-  
methylguanine, 5-methylaminomethyl uracil, 5-methoxy  
10 aminomethyl-2-thiouracil, 5-methoxyuracil, 2-methylthio-  
N<sup>6</sup>-isopentenyladenine, pseudouracil, 5-methyl-2-  
thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil,  
2-thiocytosine, and 2,6-diaminopurine. In addition to  
these base analogs, pyrimidine analogs including 6-  
15 azacytosine, 6-azathymidine and 5-trifluoromethyluracil  
described in Cook, D. P., et al, International  
Publication No. WO 92/02258 (incorporated herein by  
reference) can be conveniently incorporated into the  
invention oligomers.

20 Bases of formula (1) or (2) containing sulfur  
at the 2 and/or 4 position can be incorporated into  
oligomers and derivatized with alkynyl as R<sup>2</sup> essentially  
as described above. In corporation of 4-thiouridine and  
2-thiothymidine into oligomers has been described  
25 (Nikiforov, T. T., et al, Tet Lett (1992) 33:2379-2382;  
Clivio, P., et al Tet Lett (1992) 33:65-68; Nikiforov, T.  
T., et al, Tet Lett (1991) 32:2505-2508; Xu, Y.-Z., et al  
Tet Lett (1991) 32:2817-2820; Clivio, P., et al Tet Lett  
(1992) 33:69-72; Connolly, B.A., et al., Nucl. Acids Res.  
30 (1989) 17:4957-4974).

Preferred bases are of the formula (1) and (2)  
but also include adenine, guanine, thymine, uracil,  
35 cytosine, 5-methylcytosine, 8-oxo-N<sup>6</sup>-methyladenine,

-40-

ps ud isocytosine, and 7-deazaxanthosine. Synthesis and use of oligomers that bind to duplex DNA sequences via GT binding motif containing 7-deazaxanthosine is described in commonly owned pending U.S. application serial number 5 07/787,920, filed November 7, 1991, incorporated herein by reference.

### Synthesis

Oligomers or the segments thereof are  
10 conventionally synthesized. The synthetic methods known in the art and described herein can be used to synthesize oligomers containing bases of the invention, as well as other bases known in the art, using appropriately protected nucleomonomers (see Figure 12). Methods for  
15 the synthesis of oligomers are found, for example, in Froehler, B., et al., Nucleic Acids Res (1986) 14:5399-5467; Nucleic Acids Res (1988) 16:4831-4839; Nucleosides and Nucleotides (1987) 6:287-291; Froehler, B., Tetrahedron Letters (1986) 27:5575-5578; Caruthers, M. H.  
20 in Oligodeoxynucleotides-Antisense Inhibitions of Gene Expression (1989), J. S. Cohen, editor, CRC Press, Boca Raton, p7-24; Reese, C.B. et al, Tetrahedron Letters (1985) 26:2245-2248. Synthesis of the methylphosphonate linked oligomers via methyl phosphoramidite chemistry has  
25 also been described (Agrawal, S. et al., Tetrahedron Letters (1987) 28:3539-3542; Klem, R. E., et al, International Publication Number WO 92/07864).

Oligomers of the invention containing bases of formula (1) or (2) and one or more substitute linkages  
30 can be synthesized by one or more of four general methods according to the reaction conditions required for synthesis of a given substitute linkage. In the first  
35 method (#1), nucleomonomers containing bases of formula

-41-

(1) or (2) are directly incorporated into oligomers or a convenient fragment thereof using standard synthesis conditions and reagents. Exemplary schemes are shown in Figures 5 and 12 and exemplary linkages that can be made by method #1 include phosphodiester, phosphorothioate, phosphoroamidate, methylphosphonate, phosphorodithioate, carbonate, morpholino carbamate and sulfonate.

Method #2 involves synthesis of short synthons (dimers, trimers, etc) starting with an appropriate precursor such as a 5-bromo or 5-iodo precursor (as described below) which is subsequently converted to the C-5 substituent of formula (1) or (2) and a synthon suitable for incorporation into oligomers. This approach is exemplified in Figures 7, 8 and 11 and is suitable for synthesis of linkages including N-methylhydroxylamine, dimethylhydrazo, sulfamate, carbamate, sulfonate, sulfonamide, formacetal thioformacetal and carbonate.

Synthesis method #3 starts with uridine or cytidine (unprotected or N-protected) nucleomonomers which is subsequently iodinated. Introduction of the R<sup>2</sup> group at the 5-position is accomplished within the synthetic route to the desired dimer or trimer synthon. Method #3 is exemplified in Figure 9 and is suitable for synthesis of linkages including N-methylhydroxylamine, dimethylhydrazino, sulfamate, formacetal, thioformacetal, riboacetal, sulfonate, sulfonamide, carbamate, carbonate and boranophosphate linkages.

Method #4 starts with either (1) uracil or cytosine base containing R<sup>2</sup>, followed by conversion to a nucleomonomer suitable for incorporation into oligomers (e.g. amide linkages) as exemplified in Figure 12 or (2) a suitable precursor such as 5-iodocytosine, 5-iodouracil, cytosine or uracil which is glycosylated or alkylated

-42-

followed by conversion of the nucleomonomer to a derivative containing  $R^2$  and converted to the desired synthon (e.g. linkages such as sulfide, sulfoxide or sulfonate).

5           In general, reaction conditions that are needed to synthesize a particular dimer, trimer or larger synthon and may not be compatible with an  $R^2$  alkynyl 2'-deoxyuridine or 2'-deoxycytidine are (1) electrophilic addition reactions, or conditions that could promote  
10 electrophilic addition to C-C multiple bonds (e.g. HCl, HF,  $BF_3$ ,  $Cl_2$  or  $Br_2$ ); (2) conditions that could promote reduction of C-C multiple bonds (e.g., hydrogenation via  $H_2/Pd/C$  or hydrides such as  $B_2H_6$ ,  $BH_3$ -complex or as a class, tin hydrides or aluminum hydrides); (3) conditions  
15 that promote free radical reactions (e.g.,  $Cl_2/h\nu$ , peroxides or AIBN); and (4) reaction conditions that promote oxidation of C-C multiple bonds (e.g.  $KMnO_4$ ,  $OsO_4$ , alkyl carboxylic peracids). Synthetic schemes involving these reaction conditions may prevent the use of Method  
20 #1.

          In general, reaction conditions that are required to synthesize certain oligomers that may not be compatible with 5-iodo-2'-deoxyuridine or 5-iodo-2'-deoxycytidine, or the like, are (1) conditions that  
25 promote reduction of aryl iodides (e.g.,  $H_2$  or hydrides), (2) alkylation and arylation reactions mediated by organometallic reagents or (3) reactions that promote free radical reactions (e.g.,  $Cl_2/h\nu$ , peroxides or AIBN). Synthetic schemes involving these reactions may prevent  
30 use of Method #2.

          Method #3 starts with 2'-deoxyuridine or 2'-deoxycytidine and the nucleomonomer is subsequently converted to the 5-iodo (or 5-bromo or 5-triflate)  
35

-43-

derivative (Robins, M.J. et al, Can. J. Chem (1982) 60:554-557; Chang, P. K. et al, J Med Chem (1963) 6:428-430; Crisp, G.T., et al., Tet Lett (1990) 31:1347-1350; and Torrence, P.F., et al., J. Med. Chem (1979) 22:316-319) at the desired step, followed by conversion to a R<sup>2</sup> substituent at desired step. In some schemes it is advantageous to convert 2'-deoxyuridine to 5-R<sup>2</sup>-2'-deoxycytidine as needed by previously desired methods (Divakar, K. J., et al, J Chem. Soc. Perkin Trans I (1982) p 1171-1176). Where one of these reactions or conditions are used for synthesis of a given oligomer or fragment thereof, a nucleomonomer such as 2'-deoxyuridine can be utilized followed by conversion to the R<sup>2</sup> derivative and the cytidine derivatives thereof.

Additional exemplary linkages that can be synthesized by these general methods are summarized in Table A below.

Table A

	Linkage Structure*	Method	Reference **
20	2' -S-CH <sub>2</sub> - 5'	1-4	1
	3' -S-CH <sub>2</sub> - 5'	1-4	2
	2' -S(O)-CH <sub>2</sub> - 5'	1-4	1
	3' -S(O)-CH <sub>2</sub> - 5'	1-4	1
25	2' -S(O)(O)-CH <sub>2</sub> - 5'	1-4	1
	3' -S(O)(O)-CH <sub>2</sub> - 5'	1-4	1
	2' -CH <sub>2</sub> -S- 5'	3,4	1
	3' -CH <sub>2</sub> -S- 5'	3,4	2
	2' -CH <sub>2</sub> -S(O)- 5'	3,4	1
30	3' -CH <sub>2</sub> -S(O)- 5'	3,4	2
	2' -CH <sub>2</sub> -S(O)(O)- 5'	3,4	1
	3' -CH <sub>2</sub> -S(O)(O)- 5'	3,4	2
35	2' -CH <sub>2</sub> -CH <sub>2</sub> -O- 5'	3,4	1

-44-

	3' -CH <sub>2</sub> -CH <sub>2</sub> -O- 5'	3,4	2
	2' -N(C(O)(OR <sup>A</sup> ))-CH <sub>2</sub> -CH <sub>2</sub> - 5'	3,4	1
	3' -N(C(O)(OR <sup>A</sup> ))-CH <sub>2</sub> -CH <sub>2</sub> - 5'	3,4	2
	2' -S-CH <sub>2</sub> -CH <sub>2</sub> - 5'	3,4	1
5	3' -S-CH <sub>2</sub> -CH <sub>2</sub> - 5'	3,4	2
	2' -NH-C(O)-O- 5'	3,4	1
	2' -O-CH <sub>2</sub> -S- 5'	2-4	1
	2' -O-C(O)-N(R <sup>B</sup> )- 5'	2-4	1
	5' morpholino N-CH <sub>2</sub> - 5'	1-4	2
10	3' -X-C((CH <sub>2</sub> ) <sub>2</sub> NR <sup>C</sup> (CH <sub>2</sub> ) <sub>2</sub> )-X- 5'	2-4	3
	3' -X-C((CH <sub>2</sub> ) <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> )-X- 5'	2-4	3
15	3' -X-C((CH <sub>2</sub> ) <sub>2</sub> S(O)(O)(CH <sub>2</sub> ) <sub>2</sub> )-X- 5'	2-4	3
	3' -X-C((CH <sub>2</sub> ) <sub>2</sub> S(O)(CH <sub>2</sub> ) <sub>2</sub> )-X- 5'	2-4	3
	3' -X-C((CH <sub>2</sub> N(R <sup>C</sup> )(CH <sub>2</sub> ) <sub>2</sub> )-X- 5'	2-4	3
20	3' -X-C((CH <sub>2</sub> N(R <sup>C</sup> )(CH <sub>2</sub> N(R <sup>C</sup> ))-X- 5'	2-4	3

---

\* R<sup>A</sup> = C<sub>1-4</sub> alkyl, e.g. CH<sub>2</sub>CH<sub>3</sub> or (CH<sub>2</sub>)<sub>3</sub>CH<sub>3</sub>; R<sup>B</sup> = H or C<sub>1-4</sub> alkyl, e.g. CH<sub>3</sub>; X = O or S;

25 R<sup>C</sup> = C<sub>1-4</sub> alkyl, CN or C<sub>1-4</sub> haloalkyl, e.g. CF<sub>3</sub>; the linkages indicate covalent attachment of the indicated atom with either a 2', 3' or 5' carbon of ribose or deoxyribose.

\*\* 1 - Synthesis is accomplished essentially as described in PCT/US91/06855 for equivalent 3',5' linkages.

30 2 - International Application Number PCT/US91/06855.

35 3 - International Application Number PCT/US90/06110; linkages having a structure such as C((CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>2</sub>O) are cyclic ketals.



-45-

In addition to the substitute linkages given in Table A, Figure 17 shows a series of repeating nucleomonomer units (17-1) and exemplary amide linked oligomers (17-2, 17-3) containing selected repeating units that can contain the base analogs of the invention. In Figure 17-1,  $X^9$  is S, O, SO, SO<sub>2</sub>, CH<sub>2</sub>, CHF, CF<sub>2</sub>, or NR<sup>10</sup> and R<sup>10</sup> is (independently) H, F, OH, OCH<sub>3</sub>, CH<sub>3</sub>, or CH-lower alkyl provided that adjacent  $X^9$  are not both O. In Figures 17-2 and 17-3, each Y is independently selected and has the meaning described above (e.g. Y is H, an oligomer, a blocking group such as FMOC, tBOC, OH, DMT, MMT or an coupling group suitable for oligomer synthesis). Nucleomonomers required to synthesize oligomers containing such linkages are synthesized by method #4.

Oligomers of the invention can be synthesized by any suitable chemistry including amidite, triester or hydrogen phosphonate coupling methods and conditions. The oligomers are preferably synthesized from appropriate starting synthons such as nucleomonomers of formula (3) or (4) wherein R<sup>1</sup> at the 5'- position is DMT, MMT, FMOC (9-fluorenylmethoxycarbonyl), PACO (phenoxyacetyl), a silyl ether such as TBDMS (t-butyldiphenylsilyl) or TMS (trimethylsilyl) and R<sup>1</sup> at the 3'- position is an ester, H-phosphonate, an amidite such as  $\beta$ -cyanoethylphosphoramidite, a silyl ether such as TBDMS or TMS or t-butyldiphenyl. Alternatively, appropriate uridine or cytidine precursors such as blocked 5-iodo-2'-deoxyuridine, 5-iodo-2'-O-alkyluridine, 5-bromo-2'-deoxyuridine, 5-trifluoromethanesulfonate-2'-deoxyuridine, 5-bromo-2'-O-alkyluridine or blocked and protected 5-iodo-2'-deoxycytidine, 5-bromo-2'-deoxycytidine, 5-trifluoromethanesulfonate-2'-

-46-

deoxycytidine, 5-iodo-2'-O-alkylcytidine, 5-bromo-2'-O-alkylcytidine can be conveniently incorporated into short oligomers such as dimer, trimer, tetramer, pentamer or longer synthons that are subsequently derivatized to yield R<sup>2</sup> at the 5- position and then incorporated into suitable synthons and longer oligomers.

Synthesis of oligomers containing about 4 or more nucleomonomer residues is preferably accomplished using synthons such as monomers, dimers or trimers that carry a coupling group suitable for use with amidite, H-phosphonate or triester chemistries. The synthon can be used to link the oligomer via a phosphodiester or phosphorous-containing substitute linkage (phosphorothioate, methylphosphonate, thionomethylphosphonate, phosphoramidate and the like).

Synthesis of other nonphosphorous-containing substituted linkages can be accomplished using appropriate precursors as described herein (Figures 7-10) and are known in the art.


In addition to employing these very convenient and now most commonly used, solid phase synthesis techniques, oligomers can also be synthesized using solution phase methods such as triester synthesis. These methods are workable, but in general, less efficient for oligomers of any substantial length.

#### Intermediates or starting materials

In other aspects, the invention is directed to intermediates in the synthesis of the oligomers of the invention, including nucleomonomer analogs of formula (3) or (4):

35

-47-



15

wherein each  $R^1$  is independently H or a blocking group;  
 $R^2$  and X are as defined above;

$R^3$  is selected from the group consisting of H,  
OH, F,  $NH_2$ , OR or SR, wherein OR is O-allyl or SR is S-  
20 allyl or O or S alkyl ( $C_{1-3}$ ), wherein alkyl including  
methyl, ethyl and propyl); and

Pr is (H), or a protecting group;

provided that if X is O,  $R^3$  is H or OH, and  
both  $R^1$  are H, then  $R^2$  is 1,3-pentadiynyl, 2-, 3- and 4-  
25 pyridine-ethynyl, 2-, 4- and 5-pyrimidine-ethynyl,  
triazine-ethynyl, aryloethynyl, 2-, 4- and 5-pyrimidinyl,  
2- and 4-imidazolyl, 2- and 3-pyrrolyl-ethynyl, 2- and 3-  
furanyl-ethynyl, 2- and 3-thienyl-ethynyl, 2- and 4-  
imidazolyl-ethynyl, 2-, 4- and 5-thiazoyl-ethynyl or 2-,  
30 4- and 5-oxazolyl-ethynyl, 4-oxazolyl, 4-thiazoyl, 3-  
pyrrolyl, 3-furanyl, 3-thienyl and

further provided that for formula 3, when X is  
O, and  $R^1$  and  $R^3$  are H, then  $R^2$  is not vinyl, 1-butenyl,  
35

-48-

1-pentenyl, 1-hexenyl, 1-heptenyl, 1-octenyl, 1-propynyl, 1-butyryl, 1-hexynyl, 1-heptynyl or 1-octynyl.


Suitable protecting groups (Pr) include diisopropylformanidine, di-n-butylformanidine, diisobutylformamidine and benzoyl and suitable R<sup>1</sup> groups including DMT, MMT, FMOC, a phosphoramidite such as  $\beta$ -cyanoethylphosphoramidite, hydrogen-phosphonate and methylphosphonamidite.

Preferred protected nucleomonomers are nucleomonomers of formula (3) and (4) where X is O, R<sup>1</sup> at the 5' position is DMT, MMT or FMOC; R<sup>1</sup> at the 3' position is N,N-diisopropylamino- $\beta$ -cyanoethoxyphosphine, N,N-diisopropylaminomethoxyphosphine or hydrogen phosphonate; R<sup>2</sup> is 1-propynyl, 3-methyl-1-butyryl, 2-pyrrolyl, 2-thienyl, 2-imidazolyl or 2-thiazolyl; and Pr is (H), or diisobutylformamidine.

Preferred Embodiments:

One group of preferred oligomers of the present invention can be represented by the formula (16):

-49-



wherein each B, R<sup>1</sup> and R<sup>3</sup> are independently selected and have the meanings defined above;

25 n is an integer from 0 to 98 (values of 0 to 28 are preferred); and

each X<sup>1</sup> is independently -P(S)(O)-, -P(O)(O)- or -P(CH<sub>3</sub>)(O)-, -P(CH<sub>3</sub>)(S)-,

30 provided that at least one B is of the formula (1) or (2) as defined above; and

further provided that when at least one of said nucleomonomers of said oligomer comprises deoxyuridine 5-substituted by vinyl, 1-butenyl, 1-pentenyl, 1-hexenyl,

35

-50-

1-h ptenyl 1-octenyl, 1-propynyl, 1-butynyl, 1-hexynyl,  
1-h ptynyl, or 1-octynyl, then the remainder of the  
nucleomonomers comprising said oligomer are not solely  
comprised of phosphodiester linked 2'-deoxyadenosine, 2'-  
5 deoxyguanosine, 2'-deoxycytidine, thymidine or a  
combination thereof. Methylphosphonate,  
thionomethylphosphonate or phosphorothioate substitute  
linkages enhance the nuclease stability of the oligomers  
while their negative impact on oligomer affinity for  
10 target nucleic acids is compensated by the inclusion of  
the 5-substituted pyrimidines of the invention.

The most preferred R<sup>2</sup> group is 1-propynyl.  
Preferred R<sup>3</sup> groups are H, OH, F and O-allyl.

Other preferred oligomers of the invention  
15 contain substitute linkages other than phosphodiester,  
phosphorothioate, thionomethylphosphonate or  
methylphosphonate. Particularly useful forms of these  
substitute linkages include riboacetal, formacetal and  
3'-thioformacetal linkages, with 3'-thioformacetal being  
20 most preferred.

For synthesis of oligomers containing  
formacetal-type substitute linkages, in lieu of at least  
some phosphodiester linkages, dimeric synthons of the  
formula (6) shown in Figure 1, wherein the substituents  
25 B, X, R<sup>1</sup> and R<sup>3</sup> are as defined above are particularly  
useful.

The foregoing synthon is obtained by first  
preparing the 5-iodo pyrimidine forms of B and then  
converting these to 5-propyne derivatives, for example,  
30 by treating the dimer synthon with propyne in the  
presence of palladium, CuI, triethylamine, and DMF.  
These synthons can be incorporated into an oligomer using  
standard synthesis techniques as shown in Figures 7, 8, 9  
35

-51-

and 11. Synthesis of formacetal and 3'-thioformacetal  
substitute linkages is described in commonly owned  
pending U.S. application serial numbers 07/874,334, filed  
April 24, 1992, and 07/690,786, filed April 24, 1991,  
5 which applications are incorporated herein by reference.  
Trimer synthons containing formacetal, 3'-thioformacetal,  
riboacetal or other substitute linkages are also  
preferred compounds. Trimers and tetramers are preferred  
for synthesis of oligomers having enhanced permeation  
10 across cell membranes.

The synthesis of oligomers containing  
methylphosphonate and phosphodiester linkages is effected  
using art-known solid-phase oligomer synthesis  
techniques. A description of modifications useful in the  
15 synthesis of phosphorothioate linked oligomers are found,  
for example, in EP publication 288,163; wherein the  
oxidation step in solid phase automated synthesis using  
amidite chemistry can be independently adjusted at any  
step to obtain the phosphorothioate. An alternate method  
20 for synthesis of oligomers with phosphorothioate  
linkages, via hydrogen phosphonate chemistry, has also  
been described (Froehler, B., et al., Nucleic Acid Res  
(1986) 14:5399-5467; Froehler, B., Tetrahedron Letters  
(1986) 27:5575-5578). Sulfurization can be accomplished  
25 using reagents such as tetraethylthiuram disulfide,  
dibenzoyl tetrasulfide, thiophosphoric acid disulfide and  
the like, 3H-1,2-benzodithiol-3-one 1,1-dioxide and the  
like as described (Vu, H. et al, Tet Lett (1991) 26:3005-  
3008; Rao, M.V., et al, Tet Lett (1992) 33:4839-4842;  
30 U.S. Patent 5,151,510 issued September 29, 1992; Iyer,  
R., et al, J Org Chem (1990) 55:4693-4699; Dahl, O.  
Sulfur Reports (1991) 11:167-192). These sulfurization  
reagents can be used with either phosphoramidite or  
35

-52-

hydrogen-phosphonate chemistries. Synthesis of phosphorothioate oligomers having controlled stereochemistry can be used to generate stereoregular invention oligomers as described (International Publication No. EP 0 506 242). The thionomethyl phosphonate is prepared with methylphosphonamidite followed by sulfurization as described (Roelen, H.P.C.F., et al, Tet Lett (1992) 33:2357-2360) or with the sulfurization reagents described above.

#### Covalent Bonding Moiety

Included in some of the oligomers of the invention is a moiety which is capable of effecting at least one covalent bond between the oligomer and the target sequence. Multiple covalent bonds can also be formed by providing a multiplicity of such moieties. The covalent bond is preferably to a base residue in the target strand, but can also be made with other portions of the target, including the sugar or phosphodiester. The reaction nature of the moiety which effects crosslinking determines the nature of the target in the duplex. Preferred crosslinking moieties include acylating and alkylating agents, and, in particular, those positioned relative to the sequence specificity-conferring portion so as to permit reaction with the target location in the strand.

The crosslinking moiety can conveniently be placed as an analogous pyrimidine or purine residue in the sequence of the oligomer. The placement can be at the 5'- and/or 3'- ends, the internal portions of the sequence, or combinations of the above. Placement at the termini to permit enhanced flexibility is preferred.



-53-

Analogous moieties can also be attached to peptide backbones.

In one preferred embodiment of the invention, a switchback oligomer containing crosslinking moieties at either end can be used to bridge the strands of the duplex with at least two covalent bonds. In addition, oligomer sequences of inverted polarity can be arranged in tandem with a multiplicity of crosslinking moieties to strengthen the complex.

Exemplary of alkylating moieties that are useful in the invention include N<sup>4</sup>,N<sup>4</sup>-ethanocytosine and N<sup>6</sup>,N<sup>6</sup>-ethanoadenine.


It is clear that the base need not be a purine or pyrimidine; indeed the moiety to which the reactive function is attached need not be a base at all. Any means of attaching the reactive group is satisfactory so long as the positioning is correct.

#### Inverted Polarity

In their most general form, inverted polarity oligomers, that can incorporate one or more nucleomonomers described above, contain at least one segment along their length of the formula:



or



where -C- symbolizes any method of coupling the nucleomonomer sequences of opposite polarity (Froehler, B.C., et al Biochemistry (1992) 31:1603-1609; Horne, D.A., et al J Am Chem Soc (1990) 112:2435-2437; Beal, P.A., et al J Am Chem Soc (1992) 114:4976-4978).

-54-

In these formulas, the symbol 3'----5' indicates a stretch of oligomer in which the linkages are consistently formed between the 5'-hydroxyl of the ribosyl residue of the nucleomonomer to the left with the 3'- (or 2'- for oligomers having 2', 5' linkages) hydroxyl of the ribosyl residue of the nucleomonomer to the right (i.e., a region of uniform polarity), thus leaving the 5'-hydroxyl of the rightmost nucleomonomer ribosyl residue free for additional conjugation.

Analogously, 5'----3' indicates a stretch of oligomer in the opposite orientation wherein the linkages are formed between the 3'-hydroxyl of the ribosyl residue of the left nucleomonomer and the 5'-hydroxyl of the ribosyl residue of the nucleomonomer on the right, thus leaving the 3'-hydroxyl of the rightmost nucleomonomer ribosyl residue free for additional conjugation.

The linkage, symbolized by -C-, can be formed so as to link the 5'-hydroxyls of the adjacent ribosyl residues in formula (1) or the 3' hydroxyls of the adjacent ribosyl residues in formula (2), or the "-C-" linkage can conjugate other portions of the adjacent nucleomonomers so as to link the inverted polarity strands. "-C-" can represent a linker moiety, or simply a covalent bond.

It should be noted that if the linkage between strands of inverted polarity involves a sugar residue, either the 3'- or 2'- position can be involved in the linkage, and either of these positions can be in either R or S configuration. The choice of configuration will in part determine the geometry of the oligomer in the vicinity of the linkage. Thus, for example, if adjacent 3'- positions are used to effect a covalent linkage, less

-55-

severe deformation of the oligomer chain will generally occur if both 3'-hydroxyls involved in the linkage are in the conventional R configuration. If they are both in the S configuration, this will result in a favorable  
5 "kink" in the chain.

In addition to the use of standard oligonucleotide synthesis techniques or other couplings to effect the 5'-5' or 3'-3' linkage between ribosyl moieties, alternative approaches to joining the two  
10 strands of inverted polarity can be employed. For example, the two appended bases of the opposing termini of the inverted polarity oligomer sequences can be linked directly or through a linker, or the base of one can be linked to the sugar moiety of the other. Any suitable  
15 method of effecting the linkage can be employed. The characterizing aspect of the switchback oligomers of the invention is that they comprise tandem regions of inverted polarity, so that a region of 3'-5' polarity is followed by one of 5'-3' polarity, or vice versa, or  
20 both.

Depending on the manner of coupling the segments with inverted polarity, this coupling can be effected by insertion of a dimer wherein the appropriate 3'-positions of each member of the dimer or the 5'-  
25 positions of each member of the dimer are activated for inclusion of the dimer in the growing chain, or the conventional synthesis can be continued using the condensing nucleomonomer which is blocked in the inverse manner to that which would be employed if the polarity of  
30 the chain were to remain the same. This additional nucleomonomer can also contain a linker moiety which can be included before or after condensation to extend the chain.

35

-56-

Th synthesis of oligomers having modified residues and/or inverted polarity can be accomplished utilizing standard solid phase synthesis methods.

In general, there are two commonly used  
5 solid phase-based approaches to the synthesis of oligomers containing conventional 3'→5' or 5'→3' linkages, one involving intermediate phosphoramidites and the other involving intermediate phosphonate linkages. In the phosphoramidite based synthesis, a suitably  
10 protected nucleomonomer having a cyanoethylphosphoramidite at the position to be coupled is reacted with the free hydroxyl of a growing nucleomonomer chain derivatized to a solid support. The reaction yields a cyanoethylphosphite, which linkage must  
15 be oxidized to the cyanoethylphosphate at each intermediate step, since the reduced form is unstable to acid. The H-phosphonate-based synthesis is conducted by the reaction of a suitably protected nucleomonomer containing an H-phosphonate moiety at a position to be  
20 coupled with a solid phase-derivatized nucleomonomer chain having a free hydroxyl group, in the presence of a suitable activator to obtain an H-phosphonate diester linkage, which is stable to acid. Thus, the oxidation to the phosphate or thiophosphate can be conducted at any  
25 point during the synthesis of the oligomer or after synthesis of the oligomer is complete. The H-phosphonates can also be converted to phosphoramidate derivatives by reaction with a primary or secondary amine in the presence of carbon tetrachloride. To indicate the  
30 two approaches generically, the incoming nucleomonomer is regarded as having a "coupling phosphite/phosphate" group.

35

-57-

Variations in the type of substitut linkage are achieved by, for example, using the methyl phosphonate precursors rather than the H-phosphonates per se, using thiol derivatives of the nucleomonomer moieties and generally by methods known in the art. Nonphosphorous based linkages can also be used, such as the formacetal 3'-thioformacetal, 3'-amino and 5'-ether type linkages described above.

Thus, to obtain an oligomer segment which has a 3'-5' polarity, a nucleomonomer protected at the 5'-position and containing a coupling phosphite/phosphate group at the 3'- position is reacted with the hydroxyl at the 5'- position of a nucleomonomer coupled to a solid support through its 3'-hydroxyl. The resulting condensed oligomer is deprotected and the reaction repeated with an additional 5'-protected, 3'- phosphite/phosphate coupling nucleomonomer. Conversely, to obtain an oligomeric segment of 5'-3' polarity, a nucleomonomer protected in the 3'- position and containing a coupling phosphite/phosphate in the 5'- position is reacted with a oligomer or nucleomonomer attached to a solid support through the 5'- position, leaving the 3'- hydroxyl available to react. Similarly, after condensation of the incoming nucleomonomer, the 3'- group is deprotected and reacted with an additional 3'-protected, 5'-coupling nucleomonomer. The sequence is continued until the desired number of nucleomonomers have been added.

This oligomer chain elongation will proceed in conformance with a predetermined sequence in a series of condensations, each one of which results in the addition of another nucleomonomer. Prior to the addition of a nucleomonomer having a coupling phosphite/ phosphate, the protecting group on the solid support-bound nucleomonomer

-58-

is removed. Typically, for example, removal of the commonly-employed dimethoxytrityl (DMT) group is done by treatment with 2.5% v/v dichloroacetic acid/dichloromethane, although 1% w/v trichloroacetic acid/dichloromethane or  $\text{ZnBr}_2$ -saturated nitromethane, are also useful. Other deprotection procedures suitable for other protecting groups will be apparent to those of ordinary skill in the art. The deprotected nucleomonomer or oligomer bound to solid support is then reacted with the suitably protected nucleomonomer containing a coupling phosphite/phosphate. After each cycle the carrier bound nucleomonomer is preferably washed with anhydrous pyridine/acetonitrile (1:1, v/v), again deprotected, and the condensation reaction is completed in as many cycles as are required to form the desired number of congruent polarity internucleoside bonds which will be converted to phosphoramidates, phosphorodithioates, phosphorothioates or phosphodiester as desired.

In one embodiment, to provide the switchback linker, the incoming coupling, protected nucleomonomer is provided in the opposite polarity to the support-bound oligomers. Thus, for example, where the support-bound oligomer is 3'-5', the deprotected 5'-hydroxyl is reacted with a 3'-protected, 5'-coupling monomer, and the synthesis continued with monomers coupled at the 5'-position and protected at the 3'-position.

In another embodiment, to provide the switchback linker, a dimer synthon containing the linker element having one end which is coupled for condensation (such as a hydrogen phosphonate) to the support-bound oligomer and another end which is a protected hydroxyl group (or protected thio group) is condensed onto the

-59-

support-bound oligomer. The linked dimer is condensed and deprotected using the same conditions as those used to condense and deprotect the protected nucleomonomer hydrogen phosphonate. Subsequent extension of the  
5 oligomer chain then uses nucleomonomer residues which are coupled and protected in the opposite manner from those used to synthesize the previous portion of the chain.

One approach to this synthesis, using a linker already derivatized to two nucleomonomer residues which  
10 will be included in each portion of the strand is illustrated in Figure 5. The 5'-3' nucleomonomer portion of the strand is coupled using the 3'-DMT-5'-coupling phosphate nucleomonomers, as conventionally, to solid support. The switchback linker is derivatized to two  
15 nucleomonomer residues through their 3'- positions; the remaining 5'- positions are derivatized by the protecting group DMT in one nucleomonomer residue and a phosphonate residue in the other. The derivatized linker is coupled to the solid supported strand under standard reagent  
20 conditions and then deprotected conventionally. Further standard nucleomonomer coupling results in extension of the chain in the 3'-5' orientation.

A particularly preferred dimer synthon used to mediate the switchback in an oligomer is the O-xyloso  
25 linker (compound 2 and Formula (21) in Figure 5. The O-xyloso linker consists of two xylose-nucleomonomers (18) linked to each other by o-xylene at the 3'- position of each xylose sugar. The switchback linker synthon was synthesized using  $\alpha,\alpha'$ -orthodibromoxylene and 5'-DMT  
30 nucleomonomer (18) to give the dimer (19) as shown in Figure 5. The dimer was converted to the H-phosphonate (21) and was used in solid phase synthesis to generate  
35 oligomers. Linkers containing the bases (at position "B"

-60-

in Figure 5) thymine, 5-methylcytosine, 8-hydroxy-N<sup>6</sup>-methyladenine, pseudoisocytosine, 5-propynyluracil or cytosine are synthesized as homodimers. However, the switchback linker dimers can also be synthesized as mixed  
5 heterodimers that are separated chromatographically.

A particularly useful synthon in the preparation of oligomers containing inverted polarity is of formula (5) shown in Figure 1, wherein each Y and each B is independently chosen and have the meanings  
10 previously defined, and wherein one or both of the bases, B, can optionally be the modified base residues of formula (1) or (2) of the invention.

#### 2' Modified Oligomers

15 Included in some of the oligomers containing C-5 modified pyrimidines of the invention are modifications of the ribose or deoxyribose sugar. 2'-O-methyl-, 2'-O-ethyl- and 2'-O-allyl oligomers have been synthesized and shown to bind to single-stranded  
20 complementary nucleic acid sequences (Cotten, M., et al., Nucleic Acids Res (1990) 19:2629-2635; Blencowe, B.J., et al., Cell (1989) 59:531-539; Sproat, B.S., et al., Nucleic Acids Res (1989) 17:3373-3386; Inoue, H., et al., Nucleic Acids Res (1987) 15:6131-6148; Morisawa, H., et  
25 al., European Patent Serial No. 0339842; Chavis, C., et al., J Organic Chem (1982) 47:202-206; Sproat, B.S., et al., Nucleic Acids Res (1991) 19:733-738). The 2'-modified oligomers were reported to be relatively nuclease stable compared to unmodified controls.  
30 Synthesis of 2' fluoro nucleomonomers and their incorporation into oligomers has also been described (Codington, J.F., et al, J Org Chem (1964) 29:558-564; Fazakerley, G.V., et al, FEBS Lett (1985) 182:365-369).  
35



-61-

Synthesis of oligomer analogs containing the modified bases described herein would be based on methods described. Synthesis of oligomers containing 2'-amino nucleomonomers has been described (Pieken, W.A., et al, Science (1991) 253:314-317).

In an additional use of bases (1) and (2) in oligomers of the invention, 2'-O-allyl modified sugar forms of the nucleomonomers containing the 5-substituted bases (1) and (2) of the invention are included in the oligomer. Other 2'-O-allyl-substituted nucleomonomers can also be used at other positions in the oligomer. The 2'-O-allyl nucleomonomers can be prepared using standard methods; set forth below is a method for synthesis of the 2'-O-allyl derivatized nucleomonomers containing the invention pyrimidines through a common intermediate. Thus, for example, 5-(1-propynyl)uridine is first protected at the 5'- and 3'- positions using a tetraisopropylidisiloxane reagent, and then at the 4-oxy position using ortho-nitrophenol. The protected nucleoside is then converted to the 2'-O-allyl derivative with allyl ethyl carbonate; this useful intermediate is alternatively converted to the 2'-O-allyl-derivatized 5-(1-propynyl)uridine or the corresponding 5-(1-propynyl)cytidine. The sequence of reactions for this conversion are outlined in Figure 4.

The nucleomonomers derivatized at the 2'-position can be incorporated into oligomers in the same manner as underivatized forms.

Synthesis of 2'-thioalkyl nucleomonomers is accomplished as described in Figure 6. The protocol is useful for synthesis of 2'-thioalkyl pyrimidines via formation of an anhydro intermediate 7 that is subsequently converted to thioalkyl nucleomonomer (22).

-62-

The group designated W is defined as lower alkane (methyl, ethyl, propyl, isopropyl, butyl or isobutyl) or lower alkene including allyl. The protocol was used to synthesize 5'- DMT blocked 5-methyluridine 3'- H-phosphonate. The starting material 6 was obtained from 5-methyluridine (Markiewicz, W.T., J. Chem. Res (M) (1979) 0181-0197. Alternate blocking groups at the 5'- and 3'- positions, such as tetrahydropyran can also be utilized to obtain an equivalent starting material. The method shown in Figure 6 can thus be used to synthesize 2'-thioalkyl derivatives of the nucleomonomers containing the modified bases of the present invention in addition to synthesis of other modified pyrimidine nucleomonomers such as 2'-thioalkylcytidine, 2'-thioalkylthymidine, 2'-thioalkyl-N<sup>4</sup>-N<sup>4</sup>-ethanocytidine or 2'-thioalkyluridine. Conversion of the nucleomonomer (22) to other 5'- and 3'-derivatized compounds such as MMT,  $\beta$ -cyanoethylphosphoramidite, or methylphosphoramidite-blocked nucleomonomers can easily be accomplished using appropriate reagents.

#### Dimer and Trimer Synthons for Oligomers Containing Substitute Linkages

Oligomers containing substitute linkages that link adjacent nucleomonomer analog residues are preferably synthesized using suitably blocked dimer synthons as a starting material. For dimers wherein one or both base residues are 5-R<sup>2</sup>-U or 5-R<sup>2</sup>-C or related analogs, synthesis of a formacetal or 3'-thioformacetal-linked dimer is accomplished as described herein. An exemplary dimer containing a formacetal linkage of formula (6) shown in Figure 1, Y, X, B and R<sup>3</sup> are as defined herein.

-63-

Figures 7 and 8 show synthesis schemes for intermediates in oligomer synthesis. In both Figures, the structure U-I represents 5-iodouracil and U- $\equiv$ -CH<sub>3</sub> represents 5-(1-propynyl)uracil. Synthesis of a 3'-thioformacetal dimer or a trimer can conveniently be accomplished. As shown in Figure 7, a 5-iodouridine nucleomonomer, (25), protected at the 3'- position by esterification is first reacted with paraformaldehyde in the presence of HCl to obtain the derivatized nucleomonomer, (26), containing the substituent ClCHO<sub>2</sub>- at the 5'- position. The nucleomonomer can be esterified using, for example, a long-chain alkyl or aromatic acid, such as decyl, hexyl, benzoyl, or phenoxyacetyl. In this first step, the 3'-esterified nucleomonomer is treated with an excess of paraformaldehyde in an inert solvent at low temperature and anhydrous HCl is bubbled through the reaction mixture. The solution is conveniently dried and the solvent removed to obtain the intermediate.

The intermediate, (26), shown as the chloromethylether (ClCH<sub>2</sub>O-) at the 5'- position of the nucleoside, is then dissolved in an inert solvent. A solution of a second nucleomonomer (5-(1-propynyl)uridine derivative), 10, protected at the 5'- position, for example by DMT, and bearing an -SH substituent at the 3'- position along with a base, preferably diisopropylethylamine (DIPEA), in an inert solvent, is also prepared. The chloromethyl ether intermediate is added dropwise to this solution and the reaction mixture is stirred for several hours.

The reaction is washed with water, and the organic layer is separated and dried to obtain the dimerized product having the 3'-SCH<sub>2</sub>O- 5' linkage and protected at the 5'- and 3'- positions, as above. The

-64-

resulting dimer is deprotected at the 3'-position and then converted to the propyne derivative as shown and described in Example 1. Where the dimer is to be used in standard oligomer synthesis, it is converted to the hydrogen phosphonate using 2-chloro-4H-1,2,3-benzodioxaphosphorin-4-one (van Boom's reagent for phosphitylation (PA)). Figure 8 shows the synthesis of a 3'-thioformacetal trimer.

Synthesis of riboacetal linked dimers is shown in Figure 9. 5'-DMT, 3'-H-phosphonate dimers can be directly utilized for incorporation into oligomers by conventional methods or an appropriate precursor can be utilized as needed for conversion to trimer, tetramer or longer length oligomers.

#### Synthesis of Ethynyl Heteroaryl and Heteroaryl Derivatized Bases

Figures 14, 15 and 16 show synthetic schemes for synthesis of nucleomonomers having invention bases with ethynyl heteroaryl or heteroaryl groups at the 5 position. The nucleomonomers can be converted to blocked monomers suitable for incorporation into oligomers by conventional methods.

Synthesis of 5-phenyl-2'-deoxyuridine was accomplished as previously described using phenyltrimethylstannane (Crisp, G., et al., Tetrahedron Letters (1990) 31:1347-1350). An analogous protocol using pyridinyltrimethylstannane or pyridinyltributylstannane or the like as a starting material which is obtained from bromopyridine is used to synthesize 5-(2-pyridinyl)-2'-deoxyuridine (Example 15). Synthesis of heteroarylstannanes is conveniently accomplished as described (Bailey, T.R. Tet Lett (1986)

-65-

27:4407-4410; Jutzi, P. et al, J. Organometal Chem.  
(1983) 246:163-168; Molloy, K.C. et al, J. Organometal  
Chem. (1989) 365:61-73).

Synthesis of 5-substituted pyrimidine  
5 nucleomonomers with heteroaryl groups such as 2-thiazoyl,  
1-methyl-2-imidazolyl, 2-oxazoyl, 2-furanyl and the like  
can be accomplished using a published protocol  
(Wigerinck, P. et al., J Med Chem (1991) 34:2383-2389;  
Peters, D., et al, Nucleosides and Nucleotides (1992)  
10 11:1151-1173) followed by conversion to the corresponding  
nucleomonomer by standard methods (see Example 16). The  
5-cyano substituent is prepared as described (Inoue, T.,  
et al, Chem Pharm Bull (1978) 2:2657-2663) and also can  
be used as a starting electrophile to build heteroaryl  
15 substituted nucleomonomers as described (Wigerinck, P.,  
et al, J Med Chem (1991) 34:1767-1772).

Ethynyl heteroaryl derivatives are prepared  
from ethynyltrimethylsilane and an appropriate heteroaryl  
as described (Austin, W.B., et al, J Org Chem (1981)  
20 46:2280-2286) (see Example 14). The deprotected ethynyl  
is then introduced into the nucleomonomer by standard  
procedures (Examples 1 and 14).

#### Utility and Administration

25 As the oligomers of the invention are capable  
of significant single-stranded or double-stranded target  
nucleic acid binding activity to form duplexes, triplexes  
or other forms of stable association, these oligomers are  
useful in diagnosis and therapy of diseases that are  
30 associated with expression of one or more genes such as  
those associated with many different pathological  
conditions. Therapeutic applications can employ the  
oligomers to specifically inhibit the expression of genes  
35

-66-

(or inhibit translation of RNA sequences encoded by those genes) that are associated with either the establishment or the maintenance of a pathological condition.

Exemplary genes or RNAs encoded by those genes that can  
5 be targeted include those that encode enzymes, hormones, serum proteins, adhesion molecules, receptor molecules, cytokines, oncogenes, growth factors, and interleukins. Target genes or RNAs can be associated with any pathological condition such as those associated with  
10 inflammatory conditions, cardiovascular disorders, immune reactions, cancer, viral infections, bacterial infections and the like.

Oligomers of the present invention are suitable for both in vivo and ex vivo therapeutic applications.  
15 Indications for ex vivo uses include treatment of cells such as bone marrow or peripheral blood in conditions such as leukemia or viral infection. Target genes or RNAs encoded by those genes that can serve as targets for cancer treatments include oncogenes, such as ras, k-ras, bcl-2, c-myc, bcr, c-abl or overexpressed  
20 sequences such as mdm2, oncostatin M, IL-6 (Kaposi's sarcoma), HER-2 and translocations such as bcr/abl. Viral gene sequences or RNAs encoded by those genes such as polymerase or reverse transcriptase genes of CMV, HSV-1, HSV-2, HTLV-1, HIV-1, HIV-2, HBV, HPV, VZV, influenza  
25 virus, rhinovirus and the like are also suitable targets. Application of specifically binding oligomers can be used in conjunction with other therapeutic treatments. Other therapeutic indications for oligomers of the invention  
30 include (1) modulation of inflammatory responses by modulating expression of genes such as IL-1 receptor, IL-1, ICAM-1 or E-Selectin that play a role in mediating inflammation and (2) modulation of cellular proliferation  
35

-67-

in conditions such as arterial occlusion (restenosis) after angioplasty by modulating the expression of (a) growth or mitogenic factors such as non-muscle myosin, myc, fos, PCNA, PDGF or FGF or their receptors, or (b) cell proliferation factors such as c-myc. Other suitable extracellular proliferation factors such as TGF $\alpha$ , IL-6,  $\gamma$ INF, protein kinase C may be targeted for treatment of psoriasis or other conditions. In addition, EGF receptor, TGF $\alpha$  or MHC alleles may be targeted in autoimmune diseases.

Delivery of oligomers of the invention into cells can be enhanced by any suitable method including calcium phosphate, DMSO, glycerol or dextran transfection, electroporation or by the use of cationic anionic and/or neutral lipid compositions or liposomes by methods described (International Publication Nos. WO 90/14074, WO 91/16024, WO 91/17424, U.S. Patent 4,897,355). The oligomers can be introduced into cells by complexation with cationic lipids such as DOTMA (which can or can not form liposomes) which complex is then contacted with the cells. Suitable cationic lipids include but are not limited to N-(2,3-di(9-(Z)-octadecenylloxy))-prop-1-yl-N,N,N-trimethylammonium (DOTMA) and its salts, 1-O-oleyl-2-O-oleyl-3-dimethylaminopropyl- $\beta$ -hydroxyethylammonium and its salts and 1,2-bis(oleylloxy)-3-(trimethylammonio) propane and its salts.

Enhanced delivery of the invention oligomers can also be mediated by the use of (i) viruses such as Sendai virus (Bartzatt, R., Biotechnol Appl Biochem (1989) 11:133-135) or adenovirus (Wagner, E., et al, Proc Natl Acad Sci (1992) 89:6099-6013; (ii) polyamine or polycation conjugates using compounds such as polylysine,

-68-

pr tamine or N1, N12-bis(ethyl)spermine (Wagner, E., et al, Proc Natl Acad Sci (1991) 88:4255-4259; Zenke, M., et al, Proc Natl Acad Sci (1990) 87:3655-3659; Chank, B.K., et al, Biochem Biophys Res Commun (1988) 157:264-270; U.S. Patent 5,138,045); (iii) lipopolyamine complexes using compounds such as lipospermine (Behr, J.-P., et al, Proc Natl Acad Sci (1989) 86:6982-6986; Loeffler, J.P., et al J Neurochem (1990) 54:1812-1815); (iv) anionic, neutral or pH sensitive lipids using compounds including anionic phospholipids such as phosphatidyl glycerol, cardiolipin, phosphatidic acid or phosphatidylethanolamine (Lee, K.-D., et al, Biochim Biophys ACTA (1992) 1103:185-197; Cheddar, G., et al, Arch Biochem Biophys (1992) 294:188-192; Yoshimura, T., et al, Biochem Int (1990) 20:697-706); (v) conjugates with compounds such as transferrin or biotin or (vi) conjugates with compounds such as serum proteins (including albumin or antibodies), glycoproteins or polymers (including polyethylene glycol) that enhance pharmacokinetic properties of oligomers in a subject. As used herein, transfection refers to any method that is suitable for delivery of oligomers into cells. Any reagent such as a lipid or any agent such as a virus that can be used in transfection protocols is collectively referred to herein as a "permeation enhancing agent". Delivery of the oligomers into cells can be via cotransfection with other nucleic acids such as (i) expressable DNA fragments encoding a protein(s) or a protein fragment or (ii) translatable RNAs that encode a protein(s) or a protein fragment.

The oligomers can thus be incorporated into any suitable formulation that enhances delivery of the oligomers into cells. Suitable pharmaceutical

35



-69-

formulations also include those commonly used in applications where compounds are delivered into cells or tissues by topical administration. Compounds such as polyethylene glycol, propylene glycol, azone, nonoxonyl-  
5 9, oleic acid, DMSO, polyamines or lipopolyamines can be used in topical preparations that contain the oligomers.

The invention oligomers can be conveniently used as reagents for research or production purposes where inhibition of gene expression is desired. There  
10 are currently very few reagents available that efficiently and specifically inhibit the expression of a target gene by any mechanism. Oligomers that have been previously reported to inhibit target gene expression frequently have nonspecific effects and/or do not reduce  
15 target gene expression to very low levels (less than about 40% of uninhibited levels). The invention oligomers are, by comparison, extremely potent with  $IC_{50}$  values as low as 0.05  $\mu$ M in microinjection assays (Example 6, Table 4 below). These levels of potency  
20 permit application of the oligomers to cells with efficient inhibition of target gene expression while avoiding significant nonspecific effects on the cell. In view of this, it is clear that the invention oligomers represent a unique class of reagents that can be used to  
25 probe gene function and to probe the role of single stranded or double stranded nucleic acids.

Thus, the results as described herein provide a method of inhibiting expression of a selected protein or proteins in a subject or in cells wherein the proteins  
30 are encoded by DNA sequences and the proteins are translated from RNA sequences, comprising the steps of: introducing an oligomer of the invention into the cells; and permitting the oligomer to form a triplex with the  
35

-70-

DNA or RNA or a duplex with the DNA or RNA whereby expression of the protein or proteins is inhibited. The methods and oligomers of the present invention are suitable for modulating gene expression in both  
5 procaryotic and eucaryotic cells such as bacterial, parasite, yeast and mammalian cells.

The results described below (Example 6 and 7) demonstrate that the oligomers, when used to inhibit gene expression by an antisense mechanism, can be RNase H  
10 "competent" or RNase H "incompetent" species. Oligomers having modifications such as 2'- substitutions (2'-O-allyl and the like) or certain uncharged linkages (methylphosphonate, phosphoramidate and the like) are usually incompetent as a substrate that is recognized by  
15 and/or acted on by RNase H. RNase H competence can facilitate antisense oligomer function by degrading the target RNA in an RNA-oligomer duplex (Dagle, J.M., et al, Nucl Acids Res (1990) 18:4751-4757; Walder, J.A. et al, International Publication Number WO 89/05358). The  
20 enzyme cleaves RNA in RNA-DNA duplexes.

In order to retain RNase H competence, an oligomer requires a RNase H competent domain of three or more competent contiguous linkages located within it (Quartin, R.S., et al, Nucl Acids Res (1989) 17:7253-  
25 7262). Design of oligomers resistant to nuclease digestion will have terminal linkage, sugar and/or base modifications to effect nuclease resistance. Thus, the oligomers can be designed to have modified nucleomonomer residues at either or both the 5'- and/or 3'- ends, while  
30 having an internal RNase H competent domain.

Exemplary oligomers that retain RNase H competence would generally have uniform polarity and would comprise about 2 to about 12 nucleomonomers at the  
35

-71-

5'- end and at the 3'- end which stabilize the oligomer to nuclease degradation and about three to about 26 nucleomonomers that function as a RNase H competent domain between the RNase H incompetent 3'- and 5'- ends.

5 Variations on such an oligomer would include (1) a shorter RNase H competent domain comprising 1 or 2 RNase H competent linkages, (2) a longer RNase H incompetent domain comprising up to 15, 20 or more substitute linkages or nucleomonomers, (3) a longer RNase H  
10 competent domain comprising up to 30, 40 or more linkages, (4) oligomers with only a single RNase H incompetent domain at the 3' end or at the 5' end, or (5) oligomers having more than one RNase H competent domain. RNase H competence also applies as a consideration to  
15 oligomers having one or more regions of inverted polarity, to circular oligomers and to other types of oligomers.

Oligomers containing as few as about 8 nucleomonomers can be used to effect inhibition of target  
20 protein(s) expression by formation of duplex or triplex structures with target nucleic acid sequences. However, linear oligomers used to inhibit target protein expression via duplex or triplex formation will preferably have from about 12 to about 20 nucleomonomer  
25 residues.

Oligomers containing bases of the invention can be conveniently circularized as described (International Publication No. WO 92/19732; Kool, E.T. J Am Chem Soc (1991) 113:6265-6266; Prakash, G., et al. J Am Chem Soc  
30 (1992) 114:3523-3527). Such oligomers are suitable for binding to single-stranded or double-stranded nucleic acid targets. Circular oligomers can be of various sizes. Such oligomers in a size range of about 22-50  
35

-72-

nucle monomers can be conveniently prepared. The circular oligomers can have from about three to about six nucleomonomer residues in the loop region that separate binding domains of the oligomer as described (Prakash, G. 5 ibid). Oligomers can be enzymatically circularized through a terminal phosphate by ligase or by chemical means via linkage through the 5'- and 3'- terminal sugars and/or bases.

The oligomers can be utilized to modulate 10 target gene expression by inhibiting the interaction of nucleic acid binding proteins responsible for modulating transcription (Maher, L. J., et al, Science (1989) 245:725-730) or translation (Example 7 below). The oligomers are thus suitable as sequence-specific agents 15 that compete with nucleic acid binding proteins (including ribosomes, RNA polymerases, DNA polymerases, translational initiation factors, transcription factors that either increase or decrease transcription, protein-hormone transcription factors and the like). 20 Appropriately designed oligomers can thus be used to increase target protein synthesis through mechanisms such as binding to or near a regulatory site that transcription factors use to repress expression or by inhibiting the expression of a selected repressor protein 25 itself.

The invention oligomers can be designed to contain secondary or tertiary structures, such as pseudoknots or pseudo-half-knots (Ecker, D.J., et al, Science (1992) 257:958-961). Such structures can have a 30 more stable secondary or tertiary structure than corresponding unmodified oligomers. The enhanced stability of such structures would rely on the increased binding affinity between regions of self complementarity 35

-73-

in a single oligomer or regions of complementarity between two or more oligomers that form a given structure. Such structures can be used to mimic structures such as the HIV TAR structure in order to  
5 interfere with binding by the HIV Tat protein (a protein that binds to TAR). A similar approach can be utilized with other transcription or translation factors that recognize higher nucleic acid structures such as stems, loops, hairpins, knots and the like. Alternatively, the  
10 invention oligomers can be used to (1) disrupt or (2) bind to such structures as a method to (1) interfere with or (2) enhance the binding of proteins to nucleic acid structures.

In addition to their use in antisense or triple  
15 helix therapies, the oligomers of the invention can also be applied as therapeutic or diagnostic agents that function by direct displacement of one strand in a duplex nucleic acid. Displacement of a strand in a natural duplex such as chromosomal DNA or duplex viral DNA, RNA  
20 or hybrid DNA/RNA is possible for oligomers with a high binding affinity for their complementary target sequences. Therapeutic applications of oligomers by this method of use, referred to herein as D-looping or "D-loop therapy" has not previously been possible because the  
25 affinity of natural DNA or RNA for its complementary sequence is not great enough to efficiently displace a DNA or RNA strand in a duplex. Therapeutic efficacy of oligomers that function by D-looping would result from high affinity binding to a complementary sequence that  
30 results in modulation of the normal biological function associated with the nucleic acid target. Types of target nucleic acids include but are not limited to (i) gene sequences including exons, introns, exon/intron  
35

-74-

junctions, promoter/enhancer regions and 5' or 3' untranslated regions, (ii) regions of nucleic acids that utilize secondary structure in order to function (e.g. the HIV TAR stem-loop element or tRNAs), (iii) nucleic acids that serve structural functions such as telomeres or centromeres and (iv) any other duplex region. It is clear that oligomers can be synthesized with discrete functional domains wherein one region of an oligomer binds to a target by D-looping while an adjacent region binds a target molecule by say, forming a triple helix or binding as an aptamer to a protein. Alternatively, a D-looping oligomer can bind to each strand in a duplex by switching the strand to which the oligomer binds (i.e. by having one region of the oligomer that binds to one strand and another region that binds to the complementary strand). The controlling elements that dictate the mode of binding (i.e. triple helix or D-loop) are the sequence of the oligomer and the inherent affinity built into the oligomer. Base recognition rules in Watson-Crick duplex binding differ from those in Hoogsteen controlled triplex binding. Because of this, the oligomer base sequence can be used to dictate the type of binding rules an oligomer will utilize.

D-loop structures are formed in nature by enzyme-mediated processes (Harris, L.D. et al., J Biol Chem (1987) 262: 9285-9292) or are associated with regions where DNA replication occurs (Jacobs, H.T. et al., Nucl Acids Res (1989) 17:8949-8966). D-loops that arise from the binding of oligomers can result from a one or two step process. Direct displacement of a target strand will give rise to a D-loop by a single binding event. However, D-looping can also occur by forming a

35

-75-

trip1 helix which facilitates a strand displacement event leading to the D-loop.

Ribozymes containing bases of the invention can be designed in order to design species with altered characteristics. Ribozymes that cleave single stranded RNA or DNA (Robertson, D.L., et al Nature (1990) 344:467-468) have been described. Therapeutic applications for ribozymes have been postulated (Sarver, N. et al, Science (1990) 247:1222-1225; International Publication Number WO 91/04319). Secondary or tertiary structure necessary for ribozyme function can be affected by design of appropriate oligomer sequences. For example, ribozymes having targeting domains containing bases of the invention will have higher affinity, while maintaining base pairing specificity, for target sequences. Because of the higher affinity of the invention bases for their complementary sequences, shorter recognition domains in a ribozyme (an advantage in manufacturing) can be designed which lead to more favorable substrate turnover (an advantage in ribozyme function).

In therapeutic applications, the oligomers are utilized in a manner appropriate for treatment of a variety of conditions by inhibiting expression of appropriate target genes. For such therapy, the oligomers can be formulated for a variety of modes of administration, including systemic, topical or localized administration. Techniques and formulations generally can be found in Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, PA, latest edition. The oligomer active ingredient is generally combined with a carrier such as a diluent or excipient which can include fillers, extenders, binders, wetting agents, disintegrants, surface-active agents, or lubricants, depending on the

35

-76-

nature of the mode of administration and dosage forms. Typical dosage forms include tablets, powders, liquid preparations including suspensions, emulsions and solutions, granules, capsules and suppositories, as well  
5 as liquid preparations for injections, including liposome preparations.

For systemic administration, injection is preferred, including intramuscular, intravenous, intraperitoneal, and subcutaneous. For injection, the  
10 oligomers of the invention are formulated in liquid solutions, preferably in physiologically compatible buffers such as Hank's solution or Ringer's solution. In addition, the oligomers can be formulated in solid form and redissolved or suspended immediately prior to use.  
15 Lyophilized forms are also included. Dosages that can be used for systemic administration preferably range from about 0.01 mg/Kg to 50 mg/Kg administered once or twice per day. However, different dosing schedules can be utilized depending on (i) the potency of an individual  
20 oligomer at inhibiting the activity of its target DNA or RNA, (ii) the severity or extent of a pathological disease state associated with a given target gene, or (iii) the pharmacokinetic behavior of a given oligomer.

Systemic administration can also be by  
25 transmucosal or transdermal means, or the compounds can be administered orally. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include,  
30 for example, bile salts and fusidic acid derivatives for transmucosal administration. In addition, detergents can be used to facilitate permeation. Transmucosal administration can be through use of nasal sprays, for example,  
35



-77-

or suppositories. For oral administration, the oligomers are formulated into conventional oral administration forms such as capsules, tablets, and tonics.

5 For topical administration, the oligomers of the invention are formulated into ointments, salves, gels, or creams, as is generally known in the art. Formulation of the invention oligomers for ocular indications such as viral infections would be based on standard compositions known in the art.

10 In addition to use in therapy, the oligomers of the invention can be used as diagnostic reagents to detect the presence or absence of the target nucleic acid sequences to which they specifically bind. The enhanced binding affinity of the invention oligomers is an  
15 advantage for their use as primers and probes. Diagnostic tests can be conducted by hybridization through either double or triple helix formation which is then detected by conventional means. For example, the oligomers can be labeled using radioactive, fluorescent,  
20 or chromogenic labels and the presence of label bound to solid support detected. Alternatively, the presence of a double or triple helix can be detected by antibodies which specifically recognize these forms. Means for conducting assays using such oligomers as probes are  
25 generally known.

The use of oligomers containing the modified bases as diagnostic agents by triple helix formation is advantageous since triple helices form under mild conditions and the assays can thus be carried out without  
30 subjecting test specimens to harsh conditions. Diagnostic assays based on detection of RNA for identification of bacteria, fungi or protozoa sequences often require isolation of RNA from samples or organisms  
35

-78-

grown in the laboratory, which is laborious and time consuming, as RNA is extremely sensitive to ubiquitous nucleases.

5       The oligomer probes can also incorporate  
additional modifications such as modified sugars and/or  
substitute linkages that render the oligomer especially  
nuclease stable, and would thus be useful for assays  
conducted in the presence of cell or tissue extracts  
which normally contain nuclease activity. Oligomers  
10   containing terminal modifications often retain their  
capacity to bind to complementary sequences without loss  
of specificity (Uhlmann et al., Chemical Reviews (1990)  
90:543-584). As set forth above, the invention probes  
can also contain linkers that permit specific binding to  
15   alternate DNA strands by incorporating a linker that  
permits such binding (Froehler, B.C., et al, Biochemistry  
(1992) 31:1603-1609); Horne et al., J Am Chem Soc (1990)  
112:2435-2437).

20       Incorporation of base analogs of the present  
invention into probes that also contain covalent  
crosslinking agents has the potential to increase  
sensitivity and reduce background in diagnostic or  
detection assays. In addition, the use of crosslinking  
agents will permit novel assay modifications such as  
25   (1) the use of the crosslink to increase probe  
discrimination, (2) incorporation of a denaturing wash  
step to reduce background and (3) carrying out  
hybridization and crosslinking at or near the melting  
temperature of the hybrid to reduce secondary structure  
30   in the target and to increase probe specificity.  
Modifications of hybridization conditions have been  
previously described (Gamper et al., Nucleic Acids Res  
(1986) 14:9943).

35

-79-

Oligomers of the invention are suitable for use in diagnostic assays that employ methods wherein either the oligomer or nucleic acid to be detected are covalently attached to a solid support as described (U.S. Patent No. 4,775,619). The oligomers are also suitable for use in diagnostic assays that rely on polymerase chain reaction techniques to amplify target sequences according to described methods (European Patent Publication No. 0 393 744). Oligomers of the invention containing 5-modified pyrimidines are compatible with polymerases used in polymerase chain reaction methods such as the Taq or Vent™ polymerase. Oligomers of the invention can thus be utilized as primers in PCR protocols or triphosphate pyrimidine monomers having R<sup>2</sup> at the 5-position can be utilized as a substrate by DNA or RNA polymerases derived from thermophiles (Taq or Vent™) or other sources (E. coli, human, retrovirus, etc) to generate the oligomers of the invention in various diagnostic protocols. Synthesis of monomer triphosphates is accomplished by known methods (Otvos, L., et al, Nucl Acids Res (1987) 15:1763-1777).

The oligomers are useful as primers that are discrete sequences or as primers with a random sequence. Random sequence primers are generally about 6 or 7 nucleomonomers in length. Such primers can be used in various nucleic acid amplification protocols (PCR, ligase chain reaction, etc) or in cloning protocols. The 5-substitutions of the invention generally do not interfere with the capacity of the oligomer to function as a primer. Oligomers of the invention having 2'-modifications at sites other than the 3' terminal residue, other modifications that render the oligomer RNase H incompetent or otherwise nuclease stable can be

-80-

advantageously used as probes or primers for RNA or DNA sequences in cellular extracts or other solutions that contain nucleases. Thus, the oligomers can be used in protocols for amplifying nucleic acid in a sample by  
5 mixing the oligomer with a sample containing target nucleic acid, followed by hybridization of the oligomer with the target nucleic acid and amplifying the target nucleic acid by PCR, LCR or other suitable methods.

The oligomers derivatized to chelating agents such as EDTA, DTPA or analogs of 1,2-diaminocyclohexane acetic acid can be utilized in various in vitro diagnostic assays as described (U.S. Patent Nos. 4,772,548, 4,707,440 and 4,707,352). Alternatively,  
10 oligomers of the invention can be derivatized with crosslinking agents such as 5-(3-iodoacetamidoprop-1-yl)-2'-deoxyuridine or 5-(3-(4-bromobutyramido)prop-1-yl)-2'-deoxyuridine and used in various assay methods or kits as described (International Publication No. WO 90/14353).  
15

In addition to the foregoing uses, the ability of the oligomers to inhibit gene expression can be  
20 verified in in vitro systems by measuring the levels of expression in subject cells or in recombinant systems, by any suitable method (Graessmann, M., et al., Nucleic Acids Res (1991) 19:53-59).

25 All references cited herein are incorporated herein by reference in their entirety.

The following examples are intended to illustrate, but not to limit, the invention. Efforts have been made to insure accuracy with respect to numbers  
30 used (e.g., amounts, temperatures, etc.), but some experimental errors and deviations should be taken into account. Unless indicated otherwise, parts are parts by

35

-81-

weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Example 1

5    Synthesis of 5-(1-Propynyl)-2'-Deoxyuridine H-Phosphonate  
          Monomer and Oligomers Containing the Analog

In a 50 mL round bottom flask is placed:

- a) 708 mg (2 mmole) 5-iodo dU
- b) 10 mL anhydrous DMF
- 10    c) 76 mg (0.4 mmole) CuI
- d) 555  $\mu$ L (4 mmole) Et<sub>3</sub>N
- e) 231 mg (0.2 mmole) (Ph<sub>3</sub>P)<sub>4</sub>Pd
- f) saturate with propyne gas with stirring at room temperature (approx. 10 min.).

15           After 2 hours, more propyne gas is bubbled in and the reaction mixture is stirred overnight at room temperature. The following morning more propyne is bubbled in and stirred for an additional 2 hrs. To the reaction mixture is added Dowex ion-exchange resin (HCO<sub>3</sub>-  
20    form), 10 mL of MeOH and 10 mL of CH<sub>2</sub>Cl<sub>2</sub> and stirring continued for 1 hr. The resin is filtered off, washed with MeOH and the supernatant evaporated. Silica Gel chromatography yielded 517 mg (1.94 mmole, 97% yield) of product. See: Hobbs, J Org Chem (1989) 54:3420-3422.

25           The purified material was protected with a 5' DMT and phosphitylated as described (Marugg, J.E., et al, Tetrahedron Letters (1986) 27:2661-2664) and used in solid phase synthesis as described (Froehler, B.C., et al, U.S. Patent No. 4,959,463; Froehler, B.C., et al,  
30    Tetrahedron Letters (1986) 27:5575-5578).

          The following notation is used to represent the bases in the designated numbered oligomers in the Examples below: A, T, G, C, and U have their  
35

-82-

conventional meanings. C' = 5-methyl-2'-deoxycytidine,  
C\* = 5-(1-propynyl)-2'-deoxycytidine; U' = 5-(1-propynyl)-  
2'-deoxyuridine.

5

Example 2

Formation of Triple Helix Structures Using  
Oligomers (ON) Containing 5-Propynyl Uracil  
Residues that Bind to Duplex DNA

Three oligomers were synthesized as follows:

10

ON-1 5'TC'TC'TC'TC'TC'TTTT 3'

ON-2 5'TC'TC'TC'TC'TC'U'U'U'U'U' 3'

ON-3 5'TC'TC'TC'U'C'U'C'U'TU'TU' 3'

15

Each oligomer (ON) was hybridized with duplex  
DNA containing the target sequence 5'AGAGAGAGAGAAAAA 3'.  
Hybridization was carried out in 140 mM KCl, 5 mM MgCl<sub>2</sub>,  
5 mM Na<sub>2</sub>HPO<sub>4</sub>, pH 6.6. Thermal stability, T<sub>m</sub>, of the  
resulting triple helix formed between each oligomer and  
the target sequence was determined. The following T<sub>m</sub>  
values were obtained, ON-1 (control oligomer) was 42.1°C,  
ON-2 was 48.1°C and ON-3 was 55°C. The increased T<sub>m</sub>  
values of ON-2 and ON-3 were not expected and  
demonstrated that the triple helix formed was more stable  
than the corresponding control triple helix structure.

20

25

Example 3

Binding of Oligomers Containing 5-Propynyl  
Uracil or 5-Propynyl Cytosine  
to Single-Stranded RNA

Oligomers were synthesized as follows:

30

ON-1 5' TC'TC'TC'TC'TC'TTTT 3'

ON-3 5' TC'TC'TC'U'C'U'C'U'TU'TU' 3'

ON-4 5' TC'TC'TC'TC'TC'TTTT 3'

35

-83-

The oligomers were hybridized with a single-stranded target RNA sequence, 5' AAAAAGAGAGAGAGA 3', in 140 mM KCl, 5 mM MgCl<sub>2</sub>, 5 mM Na<sub>2</sub>HPO<sub>4</sub>, pH 6.6. The following T<sub>m</sub> values for the duplexes were obtained; ON-1 (control) was 65.5°C, ON-3 was 74.0°C and ON-4 was 73.0°C. Duplexes formed with ON-3 and ON-4 were more stable than the control oligomer. Surprisingly, ON-3 and ON-4 gave increased T<sub>m</sub> values which demonstrated that the duplex formed was more stable than the corresponding control duplex structure.

#### Example 4

##### Formation of Triple Helix Structures at Elevated pH

Triple helix formation at elevated pH was demonstrated using ON-1 as a control and ON-5, 5'U'C'U'C'U'C'U'C'U'C'U'U'U'U'U' 3'. Oligomers were hybridized with duplex target DNA, as described in Example 2 except that the buffer was at pH 7.4. T<sub>m</sub> values of the triple helix were then determined. ON-1 had a T<sub>m</sub> of 27.1 while ON-5 had a T<sub>m</sub> of 51.5. Thus, oligomers containing 5-propynyl uracil were capable of triplex formation at high pH levels, while the control oligomer formed triplex structure only at temperatures that are below physiological.

In an additional set of determinations, modified forms of ON-5 wherein the 5-substituent of the deoxyuridine was, instead of propynyl, 3-methylbutynyl (ON-5A) or 3,3-dimethyl butynyl (ON-5B), similar affects on the melting temperature of duplex and triple helices were observed. The results are shown in Table 1 below:

35

-84-

Table 1

	Duplex *		Triple-helix *		
	RNA	DNA	@ pH = 6.60		
	T <sub>m</sub> (°C)	T <sub>m</sub> (°C)	T <sub>m</sub> (°C)	ΔT <sub>m</sub> (°C)	
5	<hr/>				
	ON-1	63.0	54.5	39.6	--
	ON-5	79.0	65.5	64.8	+25.2
	ON-5A	73.5	65.5	55.9	+16.3
	ON-5B	68.5	66.0	42.5	+ 2.9
10	a) T <sub>m</sub> in 140 mM KCl/5mM Na <sub>2</sub> PO <sub>4</sub> /1mM MgCl <sub>2</sub> , pH 6.60.				

Example 5

Synthesis of 5-(3-Methyl-1-Butynyl)-2'-Deoxyuridine H-Phosphonate, Oligomers Containing the Analog and  
Formation of Triple Helix Structures Using the Oligomers  
 5-(3-Methyl-1-butynyl)-2'-deoxyuridine H-phosphonate was synthesized from 5-iododeoxyuridine essentially as described for 5-(1-propynyl)-2'-deoxyuridine H-phosphonate in Example 1, except that 5 equivalents of 3-methyl-1-butyne liquid was used in place of propyne. Silica gel purified material was then converted to the 5'-DMT, 3'-H-phosphonate monomer and used in solid phase synthesis to generate oligomers as follows (ON-1 was used as a control containing thymine and 5-methylcytosine as described in Example 2):

ON-1 5' TC'TC'TC'TC'TC'TTTTT 3'  
 ON-6 5' TC'TC'TC'U'C'U'CU'TU'TU'3'  
 ON-7 5' TC'TC'TC'TC'TC'U'U'U'U'U' 3'

Base residues designated U' correspond to 5-(3-methyl-1-butynyl)uracil. The oligomers were hybridized with duplex DNA containing the target sequence, 5' AGAGAGAGAGAAAAA 3'. Hybridization was carried out in the



-85-

buffer described in Example 2 at pH 6.2. ON-1 had a  $T_m$  of 51.0°C while the  $T_m$  of ON-6 was 55.2°C and that of ON-7 was 55.0°C.

5

Example 6

Stability of Duplex and Triplex Structures and Inhibition of Target Gene Expression

Thermal stability. Additional oligomers were tested with respect to thermal denaturation or stability (10  $T_m$ ) after hybridization with DNA or RNA targets to form triplex or duplex structures respectively. The DNA target used was an oligomer containing a self-complementary region and having a 4 nucleotide loop at the end of the hairpin. The targets were as follows:

15 DNA Duplex Target: 5' AGAGAGAGAGAAAAAGGA T T  
3' TCTCTCTCTCTTTTCCT T T

RNA Target: 5' AAAAAGAGAGAGAGA 3'

The assays for triple-helix binding were conducted in 140 mM KCL/5 mM  $\text{Na}_2\text{HPO}_4$ /5 mM  $\text{MgCl}_2$  at pH = (20 6.60 and the final concentration of all oligomers was ~2  $\mu\text{M}$ .)

ON-1, set forth above, was used as a control.

Test oligomers 8-10 contain substitutions of 5-propynyluracil for thymine and 5-propynylcytosine for (25 methylcytosine.)

ON-8 5'-TC'TC'TC'U'C'U'C'U'TU'TU' 3'

ON-9 5'-TC'TC'TC'TC'TC'TTTT 3'

ON-10 5'-U'C'U'C'U'C'U'C'U'C'U'U'U'U'U' 3'

The results obtained showed that, with respect (30 to triple-helix formation, the control ON-1 gave a  $T_m$  of 43.4°C; ON-8 gave an elevated  $T_m$  of 55.5°C; and ON-9 gave a  $T_m$  of 26.3°C. ON-8 containing U', showed an increase in  $T_m$  of 2.4°C/substitution ( $\Delta T_m @ 6.6 = +12.1^\circ\text{C}$ ) relative to (35

-86-

ON-1 and ON-9, containing C', showed a decrease in  $T_m$  of 3.4°C/substitution ( $\Delta T_m = 6.6 - 17.1^\circ\text{C}$ ) relative to ON-1. The  $T_m$  of a triple-helix, in which the third strand contains 2'-deoxycytidines, is pH dependent, and the  $T_m$  at different pH values (from 5.8 to 7.4) was used to examine the pH dependence of the complex. With all ON's the slope of the plot remained relatively constant (-18 to -20°C/pH unit).

The low  $T_m$  of ON-9, relative to ON's 1 and 8, can be explained in terms of basicity of the heterocycle. Titration of the hydrochloride salt of C' and C' showed that the pKa of the 5-propyne analog C' (3.30,  $\pm 0.05$ ) is 1.05 units less than the 5-methyl derivative C' (4.35,  $\pm 0.05$ ). The importance of protonation in triple-helix formation has been demonstrated and the results above indicate that a decrease in basicity of the cytosine nucleobase has a dramatic effect on the stability of the complex. Surprisingly, the large difference in pKa's of the cytosines (C' and C') has no significant effect on the slope of the  $T_m$  vs pH plot.

With respect to oligomer/RNA duplex formation, the control, ON-1 had a  $T_m$  of 65.5°C, ON-8 had a  $T_m$  of 74.0°C, ON-9 had a  $T_m$  of 73.0°C and ON-10 had a  $T_m$  of more than 90°C; ON-8 containing U', results in an increase in  $T_m$  of 1.7°C/substitution and ON-9, containing C', results in an increase in  $T_m$  of 1.5°C/substitution. Under these conditions ON-10, containing complete substitution with U' and C', has a  $T_m$  greater than 90°C (approx. 1.7°C/substitution) indicating that the increases in binding affinity of these substitutions are additive. These results show that the double-helix complex is greatly stabilized by substitution with both

-87-

C' and U' and, therefore, these analogs represent a new class of useful antisense ON's.

Binding assays were conducted using a combination of C' and U' in oligomers containing  
5 phosphorothioate internucleotide linkages as an additional modification. Phosphorothioate linkages render oligomers nuclease stable, but reduce binding affinity for complementary target sequences relative to unmodified phosphodiester linkages. Other phosphodiester  
10 linkage analogs known in the art such as alkylphosphonate, methylphosphonate, phosphoroamidate and triester linkages suffer from similar limitations. Unexpectedly, oligomers containing a heterocycle modification that enhances binding affinity (defined  
15 herein as a positive modification), such as U' or C', and a modification that reduces affinity (defined herein as a negative modification), were found to have improved binding (i.e. a greater binding affinity than predicted by the additive effects - positive and negative - of both  
20 modifications), relative to oligomers containing only the negative modification. Surprisingly, the propyne modification counteracts the negative binding effect of the phosphorothioate linkages to an unexpected degree. That is, for oligomers containing T and C' the  $\Delta T_m$   
25 between phosphodiester and phosphorothioate is 14°C (.7°C per substitution) while the  $\Delta T_m$  with U' and C' is 6.0°C (0.3°C per substitution). These results clearly demonstrate a synergistic effect between the negative modification (substitute linkage such as  
30 phosphorothioate) and the positive modification (base analog such as a 5-substituted pyrimidine) wherein the positive modification compensated to a degree that is more than additive with respect to binding affinity.  
35

-88-

Binding results ( $\Delta T_m$  relative to phosphodiester linkages) that wer obtained are shown in Table 2 below:

Table 2

5	ON	ON Linkage		$\Delta T_m$
		Diester	Thioate	
	ATTTTC'TTC'ATTTTTC'TTC'	54.0	40.0	-14.0
	AU'U'U'U'C'U'U'C'AU'U'U'U'U'U'C'U'U'C'	76.5	68.5	- 8.0
	AU'U'U'U'C'U'U'C'AU'U'U'U'U'U'C'U'U'C'	82.5	76.5	- 6.0

10 Additional data obtained in vitro with respect to duplex formation with target RNA corresponding to T antigen (TA<sub>g</sub>) show that the binding of the oligomer to the target is sequence-specific for the 5-substituted oligomers of the invention. The additional oligomers, 21 and 22, were prepared; ON-22 is a scrambled form of ON-21 which is designed to target the T antigen coding region as described above.

ON-21: AU'U'U'U'C'U'U'C'AU'U'U'U'U'U'C'U'U'C'

ON-22: U'U'AU'U'AU'C'U'U'C'U'U'C'U'U'U'U'C'U'

20 The oligomers were tested in phosphodiester (ON-21, ON-22) and phosphorothioate (ON-21A, ON-22A) form; and ON-21B and ON-22B the 2'-O-allyl T and C' oligomers. The results are shown in Table 3:

Table 3

25		$T_m$	$\Delta T_m$
	ON-21	76.5	
	ON-22	53.0	23.5
	ON-21A	68.0	
	ON-22A	42.0	26.0
30	ON-21B	70.0	
	ON-22B	45.0	25.0

35

-89-

The differences in  $T_m$  between the scrambled and unscrambled form are roughly the same regardless of the pyrimidine or linkage substitution used.

5                    Inhibition of target gene expression.

Additional oligomers designed to target T antigen (TA<sub>g</sub>) in a modification of the *in vivo* antisense assay described by Graessmann, M. et al. Nucleic Acids Res (1991) 19:53-59 were also modified to contain U' and/or C' as well as modified internucleoside linkages. The assay uses an expression vector for SV40 T antigen. Oligomers (ON-11 through ON-17) designed to target a sequence in the coding region were employed. The oligomers used in this assay were as follows:

- 15    ON 11: 5' ATTTTC'TTC'ATTTTTC'TTC' 3'  
      ON 12: phosphorothioate form of ON-11  
      ON 13: 5' AU'U'U'C'U'U'C'AU'U'U'U'U'C'U'U'C' 3'  
      ON 14: phosphorothioate form of ON-13  
      ON 15: phosphorothioate 5' C'U'U'C'AU'U'U'U'U'C'U'U'C' 3'  
20    ON 16: phosphorothioate 5' C'U'U'C'AU'U'U'U'U'C'U' 3'  
      ON 17: phosphorothioate 5' C'U'U'C'AU'U'U'U'U' 3'  
      ON 18: phosphorothioate 5' C'U'U'C'AU'U'AU'U'U'C'U'U'C' 3'  
      ON 19: phosphorothioate 5' C'U'U'U'C'U'U'C'U'U'AC'U'U'C' 3'

25    ON 18 and 19 were mismatch controls and have the same base composition found in ON 15 with sequence mismatches as shown in bold. The  $T_m$  of the oligomers with the complementary RNA were determined as described above. The nuclease stability of the oligomer in the cells and  
30    the ability of the oligomer to inhibit T antigen synthesis was also determined. The ability of ON 11 to ON 17 to confer RNase H sensitivity on the bound RNA was  
35    also determined and each oligomer was found to confer

-90-

RNase H sensitivity. D tails of the antisense assay protocol are described in Example 7. The results are shown in Table 4.

Table 4

5	<u>Oligomer</u>	<u>RNA T<sub>m</sub></u>	<u>S.N.</u> <sup>**</sup>	<u>TA<sub>g</sub></u>
				<u>IC<sub>50</sub></u> <sup>***</sup>
	ON-11	54.0°C	-	-
	ON-12	40.0°	+	n.s.
	ON-13	82.5°	-	2.5
10	ON-14	76.5°	+	0.05
	ON-15	71.0°	+	0.10
	ON-16	63.5°	+	0.25
	ON-17	53.5°	+	1.0
	ON-18	59.5°	+	0.50
15	ON-19	43.0°	+	-

\* T<sub>m</sub>, thermal stability of duplex determined under the same conditions as previously described at pH 6.6 in 1 mM MgCl<sub>2</sub>.

20 \*\* S.N., stability to nuclease digestion in live cells at 37°C; (-) nuclease sensitive, (+) nuclease resistant.

25 \*\*\* IC<sub>50</sub>, oligomer concentration (μM) showing 50% inhibition of TAG expression; (-), no inhibition of TAG expression detected; (n.s.), nonspecific inhibition at 25 μM.

30 As seen, substituting the phosphorothioate linkage for phosphodiester decreases the affinity of the oligomer for target RNA but enhances the nuclease stability of the oligomers in the cell. Replacement of the thymine and cytosine bases by the 5-substituted bases of the invention enhanced affinity of the oligomers. At an increased concentration of oligomer (diester, ON-13), the enhanced affinity of the oligomer led to detectable T

-91-

antigen synthesis inhibition. The phosphorothioate analog containing the modified bases is sufficiently stable and has sufficient affinity for the target RNA to effect inhibition of the synthesis of T antigen. The  
5 increasing  $IC_{50}$  value coupled with the decreasing  $T_m$  of ON 18 and ON 19 relative to ON-15 indicated that these oligomers were binding to target sequences less effectively as the number of mismatches increased. These results demonstrate sequence-specific inhibition of  
10 target gene expression by invention antisense oligomers.

In addition to inhibition of TAG synthesis, a phosphorothioate oligomer, ON 20, 5' U'U'GC'C'GU'U'U'U'-C'AU'C'AU'AU'U'U'AAU' 3', that is complementary to the  $\beta$ -galactosidase RNA, was able to inhibit  $\beta$ -galactosidase in  
15 a sequence specific manner with an  $IC_{50}$  of 0.25  $\mu$ M. ON 20A, ON 20 with T and C', did not inhibit  $\beta$ -galactosidase expression in a sequence specific manner.

#### Example 7

#### 20 Assay Method and Inhibition of Target Gene Expression

Assay Method. Antisense oligomers were evaluated for biological efficacy in vivo using a microinjection procedure. The protocol differs from  
25 previously described procedures by utilizing an additional coinjected gene which serves as an internal control for transfected gene expression (Graessman, M., et al., Nucleic Acids Res (1991) 19:53-59). Microinjections were performed using 5-10 copies per cell  
30 of pSV40 plasmid expressing the TAG gene along with varying amounts of antisense oligomers targeted to different regions to the TAG RNA. Coinjection markers such as 40 copies of plasmid per cell containing the  $\beta$ -  
35

-92-

galactosidase gene under the control of the RSV promoter or the chloramphenicol acetyl transferase gene under the control of the SV40 promoter were used. Marker genes are those which generate proteins that can readily be measured so that specificity of gene expression inhibition can be shown. The antisense oligomer does not affect the ability of the cells to continue to produce protein products of the marker gene. Suitable marker genes generate chloramphenicol acetyltransferase (CAT),  $\beta$ -galactosidase, luciferase or cell surface antigens, enzymes or markers for which assay methods are known. In control experiments without antisense oligomer, 100% of microinjected cells expressed the  $\beta$ -galactosidase protein at 4.5 h after injection while approximately 60% of microinjected cells expressed the TAg protein, as detected by dual label immunofluorescent labeling techniques. Target sequences in the TAg RNA included a coding sequence approximately 150 bases from the translation initiation AUG codon, sequences in the 5'-untranslated region and sequences at the AUG codon. Antisense oligomers from 9 to 20 bases in length were examined using concentrations of oligomers of between 5 nM and 25  $\mu$ M and the compounds were assayed at times ranging from 4.5 to 72 hours postinjection. CV1 or Rat2 cells were microinjected using conditions essentially as described (Graessman, M., et al., *ibid.*).

Target Sequence Binding and Target Gene

Inhibition. An oligomer (5' ATTTTC'TTC'ATTTTTC'TTC' 3') was systematically varied, using the phosphodiester antisense oligomer, ON-11, as a control. The phosphorothioate analog, ON-12, of the same oligomer was also prepared, but had no altered bases. The corresponding oligomer having the 5-substituted bases of



-93-

the invention universally substituted was prepared both in the phosphodiester, ON-13, and phosphor thioate form, ON-14; finally, the 2'-O-allyl-substituted form was tested as well.

5           As shown in Table 4, substituting the  
phosphorothioate linkage for phosphodiester decreases the  
affinity for target RNA but enhanced the nuclease  
stability. Analysis of the time course of inhibition of  
T antigen expression showed that ON-13 (phosphodiester  
10 linked oligomer containing U' and C') had activity that  
was detectable until 6 hours after microinjection into  
cells at a concentration of 25  $\mu$ M. By contrast, ON-14  
(phosphorothioate linked oligomer containing U' and C'  
with the same sequence as ON-13) was active for 48 hours  
15 after microinjection of 0.5  $\mu$ M oligomer into cells.

Both the 9-mer (5' C'U'U'C'AU'U'U'U' 3') and 11-mer  
(ON-17) phosphorothioates were able to inhibit T antigen  
synthesis when they contain the 5-substituted bases of  
the invention. However, the 9-mer had relatively weak  
20 sequence-specific inhibitory activity.

Also tested in the foregoing assay were an  
oligomer (5'UT) designed to bind to the 5' untranslated  
region of the T antigen near the CAP site of the mRNA and  
an oligomer with a sequence designed to bind to the  
25 region of the start codon. These oligomers have the  
sequences shown:

5'UT oligomer: 5'-GCC TCC TCA CTA CTT CTG GA-3'

AUG oligomer: 5'-CAT CTT TGC AAA GCT TTT TG-3'

30           The phosphorothioate form of the 5'UT and AUG  
oligomers composed of thymine and 5-methylcytosine bases  
was unable to effect detectable inhibition at 20  $\mu$ M.  
35 However, the phosphorothioate analogs wherein 5-

-94-

propynyluracil was substituted systematically for thymine residues showed 100% inhibition at 1  $\mu$ M.

Further experiments with this assay system using a T antigen target sequence in the 5' untranslated region demonstrated that substitution of the modified oligomers of the invention containing phosphodiester linkages but containing 2'-O-allyl substitutions in the oligomers containing fully substituted nucleomonomers wherein C' replaces C and U' replaces T are capable of inhibiting T antigen synthesis. Table 5 shows the results obtained using the 5'UT oligomer. Oligomers 1-4 were 20-mers having the sequence shown above. Oligomer 5 had the underlined sequence shown above and 5-propynyluridine substituted for thymidine and 5-propynylcytidine substituted for cytidine.

Table 5

	<u>Oligomer*</u>	<u>T<sub>m</sub></u>	<u>IC<sub>50</sub> (<math>\mu</math>M)</u>
	1. 2'-O-Me (U, C)	-	n.i.**
20	2. Thioate (T, C')	70.5	2.5
	3. Thioate (U', C')	81.0	0.5
	4. Thioate (U', C')	>90.0	0.25
	5. 2'-O-allyl (U', C')	>90.0	5.0

\* Thioate or phosphorothioate linkages

25 \*\* No detectable inhibition at 5  $\mu$ M

As shown in Table 5 various combinations of inclusion of U' and C' along with either a phosphorothioate backbone or a 2'-O-allyl substitution provided inhibition. Although oligomer 5 is not a substrate for RNase H, inhibition of TAG expression was observed. Inhibition mediated by oligomer 5 is believed to result from the high affinity and nuclease stability

-95-

that the 2'-O-allyl modification confers. Incorporation of U' and/or C' into oligomers containing full or partial 2'-O-allyl modification will provide oligomers that can be used to inhibit target gene expression.

5           In addition to sequences in the 5'UT, AUG codon region and exon described above, TAg sequences at TAg intron/exon junction, exon/exon junction and in an intron were targeted using 15-mer phosphorothioate oligomers that were fully substituted with U' and/or C' according to  
10           the target sequences. The oligomers contained from about 50% to about 70% of U' and/or C' bases in the oligomer. All of the oligomers effectively inhibited TAg synthesis. These results indicated that the high affinity oligomers were capable of inhibiting gene expression at locations  
15           throughout the RNA and were thus effective in spite of any secondary structure that can have been present in the TAg RNA.

#### Example 8

##### 20           Delivery of Oligomers into Cells

          ON-15 was tested for inhibition of T antigen (TAg) expression using the method described in Example 7. ON-15 was incubated for 24 hours with CV1 cells at an extracellular concentration of 50  $\mu$ M in tissue culture  
25           medium, followed by microinjection of TAg and  $\beta$ -galactosidase expression plasmids. 4.5 hours after injection, cells were fixed and stained for TAg expression. By comparison with microinjected ON-15, which efficiently inhibits TAg expression, ON-15  
30           incubated in extracellular medium was much less efficient with no detectable inhibition of TAg synthesis. The experiment was repeated using ON-15 derivatized at the 5' end with fluorescein-aminohexanol (Fl-ON-15) at an  
35

-96-

extracellular concentration of 50  $\mu$ M which was incubated with cells for 24 hours. Control cells were microinjected with Fl-ON-15 at an intracellular concentration of 0.5  $\mu$ M along with TAg and  $\beta$ -galactosidase expression plasmids. Microinjected Fl-ON-15 was localized in the nucleus while Fl-ON-15 added to the extracellular medium was localized in cytoplasmic compartments that resembled endosomes and lysosomes. The same pattern of results were obtained in CV1, Rat2, HeLa, SKOV-3, BUD-8, BC3H1 and ccd45sk cell lines. Oligomers of the invention are active in different mammalian species and are thus suitable for modulating gene expression regardless of the species.

Fl-ON-15 and ON-15 were delivered to cellular cytoplasm using a commercially available cationic lipid preparation, Lipofectin™ (BRL-Gibco, cat. no. 8292SA). A Lipofectin™ concentration of 10  $\mu$ M in Optimem (BRL-Gibco) was used according to manufacturer's instructions. DOTMA is the lipid present in Lipofectin™. Cells were incubated in Optimem containing either the Fl-ON-15-lipid or ON-15-lipid preparation for 4 hours, followed by incubation of the cells for 16 hours in standard medium (DMEM with 10% FBS for CV1 cells). Immunofluorescence analysis of treated cells showed that about 90% of the cells contained Fl-ON-15 localized in the nucleus. Delivery of ON-15 to cells was assayed by microinjection of TAg and  $\beta$ -galactosidase expression plasmids into cells after incubation with the lipid-oligomer complex. ON-15 inhibited TAg expression with an  $IC_{50}$  of less than 5 nM using Lipofectin™ at a 10  $\mu$ M concentration. Preparations of oligomer complexed with cationic lipid (such as Lipofectin™) were thus capable of delivering oligomers containing a label such as fluorescein and/or 5-modified

-97-

pyrimidines such as U' or C' to the cellular cytoplasm, indicating that modifications incorporated into oligomers, such as base analogs or a label, do not interfere with formation cationic lipid-oligomer  
5 complexes.

In an alternative protocol, Fl-ON-15 was also delivered into cells by a transfection protocol using DMSO. CV1 cells were incubated in DMEM-10% FBS medium containing 1% DMSO and 1  $\mu$ M Fl-ON-15 for 4 hours at 37°C.  
10 Fluorescence microscopy demonstrated that the oligomer was localized in the nucleus of about 20% of the treated cells.

Fl-ON-15 was synthesized by coupling a commercially available aminohexane amidite (Glen  
15 Research) to the 5'-OH of ON-15 using standard coupling conditions. The free amine was then linked to fluorescein-NHS ester (Molecular Probes) to generate Fl-ON-15. Synthesis of fluorescein-linked oligomer can also be accomplished using fluorescein amidite or fluorescein-  
20 CPG according to manufacturer's instructions (Glen Research).

#### Example 9

##### Synthesis of 2'-O-Allyl Monomers for Oligomer Synthesis

25

Preparation of 5-propynyl-2'-O-allyluridine nucleomonomer. 343 mg (0.50 mmole) of 14 (Figure 4) was dissolved into anhydrous CH<sub>3</sub>CN (5 mL) and to this was added 2-pyridinealdoxime (67 mg, 0.55 mmole) and 1,1,3,3-tetramethylguanidine (75  $\mu$ L, 0.6 mmole) at room  
30 temperature. After 18 hr the reaction mixture was diluted with EtOAc and washed with aq. citric acid (0.1 M). The aqueous layer was extracted with EtOAc, the  
35

-98-

combined organic layers washed with saturated aq. NaHCO<sub>3</sub>,  
(3 times), dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. The residue  
was dissolved into EtOAc (5 mL) and to this was added 1 M  
TBAF/THF (1.5 mL, 1.5 mmole), the solution stirred for 1  
5 hr and diluted with EtOAc. The solution was washed with  
saturated aq. NaHCO<sub>3</sub> (2 times), the combined aqueous  
layer extracted with EtOAc (3 times), the combined  
organic phase dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated. The  
residue was evaporated from anhydrous pyridine (10 mL),  
10 dissolved into anhydrous pyridine (5 mL), and to this was  
added dimethoxytrityl chloride (200 mg, 0.6 mmole) and  
the solution stirred for 18 hr. The reaction mixture was  
evaporated to approximately 2 mL, diluted with CH<sub>2</sub>Cl<sub>2</sub>,  
washed with saturated aq. NaHCO<sub>3</sub>, dried over Na<sub>2</sub>SO<sub>4</sub> and  
15 evaporated. Purification by silica gel chromatography  
(EtOAc/Hexane, 1/1) yielded 197 mg (0.32 mmole, 64%) of 6  
shown in Figure 4.

Preparation of 5-propynyl-2'-O-allylcytidine  
nucleomonomer. 343 mg (0.50 mmole) of 14 was dissolved  
20 into anhydrous CH<sub>3</sub>CN (10 mL), and the solution  
transferred to a Parr Bomb, cooled to 0°C, and saturated  
with NH<sub>3</sub>. This was placed in an 80°C bath for 24 hr (75  
psi), cooled to room temperature and evaporated to  
dryness. The residue was evaporated from anhydrous DMF  
25 (10 mL), dissolved into anhydrous DMF (5 mL), and to this  
was added diisobutylformamide dimethylacetal (0.2 mL,  
0.84 mmole) at room temperature. After 18 hr H<sub>2</sub>O (25 µL)  
was added, the solution evaporated, dissolved into EtOAc  
(5 mL) and to this was added 1 M TBAF/THF (1.5 mL, 1.5  
30 mmole). After 1 hr the reaction mixture was diluted with  
EtOAc, washed with saturated aq. NaHCO<sub>3</sub>, dried over Na<sub>2</sub>SO<sub>4</sub>  
and evaporated. The residue was evaporated from  
35 anhydrous pyridine (10 mL), dissolved into anhydrous

-99-

pyridin (5 mL), and to this was added dimethoxytrityl chloride (200 mg, 0.6 mmole) and the solution stirred for 5 hr. The reaction mixture was evaporated to approximately 2 mL, diluted with  $\text{CH}_2\text{Cl}_2$ , washed with saturated aq.  $\text{NaHCO}_3$ , dried over  $\text{Na}_2\text{SO}_4$  and evaporated. Purification by silica gel chromatography ( $\text{EtOAc/Hexane}$ , from 2/3 to 3/2) yielded 242 mg (0.32 mmole, 64%) of **8** shown in Figure 4.

10

Example 10

Formacetal Dimer Synthesis

Figure 11 shows a synthesis scheme that was used to obtain a formacetal linked 5-propynyl-2'-deoxyuracil dimer. The synthesis protocol introduced the propynyl substituent at the level of a dimer by conversion of the 5-iodouracil precursor as shown. A similar protocol can be used to convert trimers, tetramers, pentamers or longer 5-iodo precursors to the 5-propynyl product in a similar fashion. This synthetic method gave unexpectedly high yields of the 5-propynyl product.

20

Example 11

3'-Thioformacetal Dimer Synthesis

Preparation of (26): (25) was suspended into  $\text{CH}_2\text{Cl}_2$  and paraformaldehyde (1.6 eq) was added, the suspension cooled to  $0^\circ\text{C}$  and  $\text{HCl}$  (anhydrous) passed through the solution for about 10 minutes. The suspension was sealed and stored at  $0-5^\circ\text{C}$  for 4 hours. The resulting solution was dried, filtered, and evaporated to yield (26).

30

Preparation of 10: 3'-Deoxy-3'-thioacetyl-5'-dimethoxytrityl-5-propynyl-2'-deoxyuridine was dissolved in methanolic ammonia in a flask that had been flushed with  $\text{O}_2$  and the solution was sealed and stirred for 1

35

-100-

hour. The solvent was removed and the residue dissolved in EtOAc and washed with NaHCO<sub>3</sub> and brine. The organic phase was dried, evaporated and purified by silica gel chromatography (MeOH/CH<sub>2</sub>Cl<sub>2</sub>). The resulting disulfide was dissolved into dioxane/H<sub>2</sub>O followed by addition of tributylphosphine (1.0 eq) and the solution stirred for 30 minutes. The solvent was removed and crude compound 10 was used directly to prepare (27).

Preparation of (27): Compound 10 (1.1 eq) was dissolved in DMF and DIPEA (diisopropylethylamine, 2.5 eq) was added and the solution was cooled to 0°C under Ar. A solution of compound (26) (1 eq, in DMF) was added, the solution stirred for 10 hours, diluted, extracted against H<sub>2</sub>O, dried, evaporated and purified by silica gel chromatography (MeOH/CH<sub>2</sub>Cl<sub>2</sub>).

Preparation of 11: (27) was dissolved in methanolic ammonia and stirred in a sealed flask at room temperature for 3 hours. The solvent was removed and purified (Silica Gel, MeOH/CH<sub>2</sub>Cl<sub>2</sub> (0-4% MeOH)) and the propyne moiety introduced has described in Example 1.

Preparation of 11A: 11 was phosphitylated using standard procedures.

#### Example 12

#### 5-(2-Pyridinyl)-2'-Deoxyuridine and 5-(2-Pyridinyl)-2'-Deoxycytidine Synthesis

2-Trimethylstannylpyridine. A solution of 15.7 mL of 1.6 M n-butyllithium (25.1 mmol) in anhydrous ether (25 mL) was stirred under Ar at -78°C and to this was added a solution of 2-bromopyridine (3.97 g, 25.1 mmol) in anhydrous ether (12.5 mL). The resulting orange solution was stirred for 2 h, and trimethyltin chloride in THF (26.0 mmol, 26 and, 1.0 M) added over 30 min. The



-101-

5 reacti n was stirred for 1 h at -78°C then warmed to room temperature over 1 h, filtered and the filtrate evaporated. Distillation afforded 3.1 g (51%) of the title compound as a colorless liquid which solidified in the receiver flask. B.p. 65°C/2 mm Hg; Literature B.p. 75°C/4 mm Hg.

10 5-(2-pyridinyl)-2'-deoxyuridine. In a 25 mL pear shaped flask was placed 5-iodo-2'-deoxyuridine (0.354 g, 1.0 mmol), 2-trimethylstannylpyridine (0.84 g, 3.5 mmol), Bis (triphenylphosphine)-palladium (II) chloride (0.070 g, 0.1 mmol), and anhydrous 1,4-dioxane (15 mL). The reaction was heated at 60°C for 15 hrs then at 90°C for 1 h. The solvent was evaporated and the residue purified  
15 by silica gel chromatography (10% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>, (1% NH<sub>3</sub>)) to yield 0.253 g (83%) of the title compound as a white solid m.p. 201-203°C.

20 5-(2-pyridinyl)-2'-deoxycytidine. In a 25 mL pear shaped flask was placed 5-iodo-2'-deoxycytidine (0.425 g, 1.2 mmol), 2-trimethylstannylpyridine (1.4 g, 5.8 mmol), Bis (triphenylphosphine)-palladium (II) chloride (0.084 g, 0.12 mmol), and anhydrous 1,4-dioxane (15 mL). The reaction was heated at 60°C for 15 hrs then at 90°C for 1  
25 h. The solvent was evaporated and the residue purified by silica gel chromatography (10% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>, (1% NH<sub>3</sub>)) to yield 0.173 g (47%) of the title compound as a white solid m.p. 197-198°C.

30 Example 13

Synthesis of 5-(Thienyl)-5'-Dimethoxytrityl-2'Deoxyuridine  
2-Trimethylstannylthiophene. To a solution of thiophene  
35 (2.1g, 25.0 mmol) in anhydrous THF (50 mL) was added

-102-

dropwise over 30 minutes at -38 C t-butyllithium in pentane (1.6 M, 16.0 mL, 25.6 mmol). The solution was stirred for 1 h at -38 C and then cooled to -78 C. Trimethyltin chloride in THF (1 M, 25.3 mL, 25.3 mmol) was added dropwise over 30 min. The reaction was then stirred for 2 h at -78 C and then for 15 h at r.t. The solvent was evaporated and the resulting yellow solid was taken up in ether, washed with water and dried Na<sub>2</sub>SO<sub>4</sub> and then evaporated to give a light brown liquid which solidified on standing.

5-(2-thienyl)-5'-dimethoxytrityl-2'-deoxyuridine. In a 50 mL flask was placed 5-iodo-dimethoxytrityl-3'-O-toluolyl-2'-deoxyuridine (2.15g, 2.77 mmol), 2-trimethylstannylthiophene (1.85 g, 7.5 mmol), Bis(triphenylphosphine)-palladium (II) chloride (0.196 g, 0.28 mmol) and anhydrous THF (30.0 mL). The was heated at 73 C for 18 h, then cooled to room temperature. The resulting black precipitate was filtered and washed with THF. The solvent was evaporated and the green residue was dissolved in ethyl acetate (100 mL), washed with water (50 mL) and dried (Na<sub>2</sub>SO<sub>4</sub>). The solvent was evaporated and the residue was taken up in dioxane/conc. NH<sub>4</sub>OH (1/1, 80 mL), stirred for 18 h at r.t. and evaporated. Silica gel purification yielded 0.430 g (25%) of 5-(2-thienyl)-5'-dimethoxytrityl-2'-deoxyuridine.

#### Example 14

#### Synthesis of 5-((1-Ethynyl)-2-Pyrimidinyl)-2'-Deoxyuridine

2-Iodopyrimidine. Liquid HI (28.0 mL, 212 mmol) was stirred vigorously at 0°C and solid 2-chloropyrimidine (7.0 g, 61 mmol) was added slowly so that the temperature

-103-

did not rise above 8°C. The reaction was stirred at 5°C for 1 h., solid potassium carbonate (14.7g, 106 mmol) was carefully added over 0.5 h. (temperature less than 10°C), and the reaction was decolorized by addition of a small amount of solid sodium bisulfite. The solution was extracted with ether (5x50 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and the residue was recrystallized from hexane to afford 7.11g (56%) of the title compound, m.p. 29-30°C.

2-((Trimethylsilyl)ethynyl)pyrimidine. In a 50 mL flask was placed 2-iodopyrimidine (20.g, 9.7 mmol), copper (I) iodide (9.5mg, 0.05 mmol), triethylamine (0.7 mL, 5.0 mmol), Bis(triphenylphosphine)-palladium (II) chloride (0.07g, 0.1 mmol) and anhydrous dioxane (10 mL). The reaction was heated at 60°C for 18 hours then at 95°C for 8 h. The solvent was evaporated and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (200 mL), washed with H<sub>2</sub>O (40 mL), dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The product was purified by silica gel chromatography (CH<sub>2</sub>Cl<sub>2</sub>) to yield 1.06g (60%) of the title compound.

2-ethynylpyrimidine. To a solution of 2-((Trimethylsilyl)ethynyl)pyrimidine (1.06g, 6.0 mmol) in anhydrous methanol (9 mL) was added solid potassium carbonate (0.083g, 0.6 mmol) and the reaction stirred for 2.5 h. The solid was removed by filtration, washed with methanol (10 mL) and the supernatant evaporated. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (30 mL), washed with saturated NaHCO<sub>3</sub>, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. Silica gel chromatography (1-2% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>) yielded 0.400g (64%) of the title compound. This was coupled to 5-Iodo-2'-deoxyuridine and converted to the protected H-phosphonate as in Example 1.

35

-104-

Example 15

Binding of Oligomers Containing U' and 3'-Thioformacetal  
or Formacetal Linkages

The following oligomers were synthesized.

5 Unless otherwise indicated, all linkages were  
phosphodiester. The symbol, •, indicates a 3'-  
thioformacetal linkage (3',5'), and the symbol, ○, indicates a formacetal linkage (3',5').

ON-1 5' TC'TC'TC'TC'TC'TTTT 3'  
10 ON-23 5' TC'TC'TC'TC'TC'U'U'U'U'T 3'  
ON-24 5' TC'TC'TC'TC'TC'T•TT•TT 3'  
ON-25 5' TC'TC'TC'TC'TC'U'•U'•U'T 3'  
ON-26 5' TC'TC'TC'TC'TC'U'OU'OU'T 3'

The oligomers were tested with respect to  
15 thermal denaturation ( $T_m$ ) or stability after  
hybridization with DNA or RNA targets to form triplex or  
duplex structures respectively. The DNA target used was  
an oligomer containing a self-complementary region and  
having a 4 nucleotide loop at the end of the hairpin as  
20 shown. The targets were as follows:

DNA Duplex Target: 5' AGAGAGAGAGAAAAGGA T T  
3' TCTCTCTCTCTTTTCCT T T

RNA Target: 5' AAAAAGAGAGAGAGA 3'

The assays for binding were conducted in 140 mM  
25 KCL/5 mM  $\text{Na}_2\text{HPO}_4$ /1 mM  $\text{MgCl}_2$  at pH = 7.2 for single  
stranded RNA or DNA (duplex hybridization) and at pH =  
6.6 for duplex DNA (triplex hybridization conditions).  
The final concentration of all oligomers was  $\sim 2 \mu\text{M}$ .  $T_m$   
data obtained with the oligomers is shown in Table 6.

35

-105-

Table 6  
Duplex T<sub>m</sub>

ON	RNA	DNA	Triplex T <sub>m</sub>
5 ON-1	62.5	55.5	38.9
ON-23	69.0	59.5	45.0
ON-24	62.0	53.0	41.5
ON-25	71.5	61.5	47.6
10 ON-26	69.0	59.5	45.5

Example 16

Binding of Oligomers Containing 5-(2-Pyridinyl)-2'-Deoxyuridine (U<sup>p</sup>) or 5-(2-Pyridinyl)-2'-Deoxycytidine (C<sup>p</sup>) and 5-(2-Thienyl)-2'-Deoxyuridine (U<sup>t</sup>)

The following oligomers were synthesized. All linkages were phosphodiester.

ON-1 5' TC'TC'TC'TC'TC'TTTTT 3'  
 20 ON-28 5' TC'TC'TC'TC'TC'U<sup>p</sup>U<sup>p</sup>U<sup>p</sup>U<sup>p</sup> 3'  
 ON-29 5' TC'TC'TC'U<sup>p</sup>C'U<sup>p</sup>C'U<sup>p</sup>TU<sup>p</sup>TU<sup>p</sup> 3'  
 ON-30 5' TC<sup>p</sup>TC<sup>p</sup>TC<sup>p</sup>TC<sup>p</sup>TC<sup>p</sup>TTTTT 3'  
 ON-43 5' TC'TC'TC'TC'TC'U<sup>t</sup>U<sup>t</sup>U<sup>t</sup>U<sup>t</sup> 3'  
 ON-44 5' TC'TC'TC'U<sup>t</sup>C'U<sup>t</sup>C'U<sup>t</sup>TU<sup>t</sup>TU<sup>t</sup> 3'

25 The oligomers were tested with respect to thermal denaturation (T<sub>m</sub>) or stability after hybridization with ssDNA or ssRNA targets to form a duplex and with dsDNA to form a triplex. The DNA target used was an oligomer containing a self-complementary region and having a 4 nucleotide loop at the end of the hairpin as shown. The targets were as follows:

30 DNA Duplex Target: 5' AGAGAGAGAGAAAAAGGA T T  
 35 3' TCTCTCTCTCTTTTCCT T T

-106-

DNA/RNA Target Sequence: 5' AAAAAGAGAGAGAGA 3'

The assays for triple-helix binding were conducted in 140 mM KCL/5 mM Na<sub>2</sub>HPO<sub>4</sub>/1 mM MgCL<sub>2</sub> at pH = 7.2 for single stranded RNA or DNA (duplex hybridization) and at pH = 6.6 for duplex DNA (triplex hybridization conditions). The final concentration of all oligomers was ~2 μM. T<sub>m</sub> data obtained with the oligomers is shown in Table 7.

Table 7			
Duplex T <sub>m</sub>			
ON	RNA	DNA	Triplex T <sub>m</sub>
ON-1	62.5	55.0	39.6
ON-28	64.0	54.0	41.8
ON-29	62.5	52.0	31.7
ON-30	61.5	46.5	--
ON-43	64.5	55.0	31.6
ON-44	63.5	52.5	19.9

#### Example 17

Binding of Oligomers Containing 5-(1-Propynyl)-2'-Deoxyuridine (U') and Carbocyclic 5-Methyl-2'-Deoxycytidine(C') or 8-Oxo-N<sup>6</sup>-Methyl-2'-Deoxyadenosine (M)

The following oligomers were synthesized. All linkages were phosphodiester.

ON-1 5' TC'TC'TC'TC'TC'TTTTT 3'  
ON-32 5' U'C'U'C'U'C'U'C'U'C'U'U'U'U'U' 3'  
ON-33 5' TTTMTTTMMTMMTTTTT 3'  
ON-34 5' U'U'U'MU'U'U'MMU'MMU'U'U'U'U' 3'

-107-

The oligomers were tested with respect to target sequence binding by footprint analysis using DNase digestion after binding with duplex DNA to form triplex structures. Hybridization was conducted at 37 C for about 1 hour in 140 mM KCL, 10 mM NaCl, 1 mM MgCl<sub>2</sub>, 1 mM spermine hydrochloride, MOPS pH 7.2 using target DNA at approximately 1 nM. The target sequence was as follows:

DNA Duplex Target: 5' AGAGAGAGAGAAAA 3'  
3' TCTCTCTCTCTTTT 5'.

The sequence was contained on a gel purified 370-bp restriction fragment derived from a cloning vector. The concentration of ON-1 and ON-32 was varied from 0.01 to 10.0  $\mu$ M. The nuclease protection results obtained with ON-1 and ON-32 indicated that ON-32 had a binding affinity for duplex DNA that was about 1000-fold greater than the affinity of ON-1 for the same target.

The ON-33 and ON-34 target sequence was as follows:

DNA Duplex Target: 5' AAAGAAAGGAGGAAAA 3'  
3' TTTCTTTCCTCCTTTT 5'.

The sequence was contained in a plasmid that was linearized by restriction enzyme digestion. The sequence is found in the promoter region of the IL-1 gene. The assays for triple-helix binding were conducted in 140 mM KCL/10 mM NaCl/1 mM MgCl<sub>2</sub>/1 mM spermine hydrochloride/20 mM MOPS pH 7.2. The final concentration of the target was 2 nM while the concentration of ON-33 and ON-34 was varied from 0.1 to 10.0  $\mu$ M. The nuclease protection results obtained with ON-33, which gave full DNase protection at 0.1  $\mu$ M, and ON-34, which gave full DNase protection at a concentration well below 0.1  $\mu$ M, indicated that ON-34 had a binding affinity for duplex

35

-108-

DNA that was greater than or equal to 10-fold greater than the affinity of ON-33 for the same target.

The results obtained with these oligomers indicate that the invention bases are compatible with other triplex binding competent base analogs and can thus be used to design high affinity oligomers with triplex-competent bases and base analogs as desired.

Example 18

Duplex Formation Using Oligomers Containing 5-(1-Propynyl)-2'-O-Allyluridine (U<sup>x</sup>) or 5-(1-Propynyl)-2'-O-Allylcytidine (C<sup>x</sup>)

The following oligomers were synthesized. All linkages were phosphodiester. Nucleomonomers were prepared as described in Example 9. 5-Methyl-2'-O-allyluridine, T', and 5-methyl-2'-O-allylcytidine, C'', were used in addition to U<sup>x</sup> and C<sup>x</sup>.

ON-1     5' TC'TC'TC'TC'TC'TTTTT 3'  
ON-35    5' TC'TC'TC'TC'TC'T'T'T'T'T' 3'  
ON-36    5' TC'TC'TC'TC'TC'U'U'U'U'U' 3'  
ON-37    5' TC'TC'TC'TC'TC'U<sup>x</sup>U<sup>x</sup>U<sup>x</sup>U<sup>x</sup> 3'  
ON-38    5' TC''TC''TC''TC''TC''TTTTT 3'  
ON-39    5' TC'TC'TC'TC'TC'TTTTT 3'  
ON-40    5' TC<sup>x</sup>TC<sup>x</sup>TC<sup>x</sup>TC<sup>x</sup>TC<sup>x</sup>TTTTT 3'

The oligomers were tested with respect to thermal denaturation (T<sub>m</sub>) or stability after hybridization with an RNA target to form duplex structures. The target was as follows:

RNA Target:            5' AAAAAGAGAGAGAGA 3'

The assays for duplex-helix binding was conducted in 140 mM KCL/5 mM Na<sub>2</sub>HPO<sub>4</sub>/1 mM MgCL<sub>2</sub> at pH = 6.6 for the RNA target. The final concentration of all



-109-

oligomers was  $\sim 2 \mu\text{M}$ .  $T_m$  data obtained with the oligomers is shown in Table 8.

Table 8

5	ON	$T_m$	$\Delta T_m$ ( $^{\circ}\text{C}/\text{substitution}$ )
	ON-1	63.0	-
	ON-35	64.5	+0.3
10	ON-36	70.5	+1.5
	ON-37	71.5	+1.7
	ON-38	66.5	+0.7
	ON-39	70.0	+1.4
	ON-40	73.0	+2.0

15

The results obtained from these analyses demonstrated that the invention bases can be combined with modified sugars in binding competent oligomers. Enhanced binding due to the 2'-O-allyl sugar modification and to the invention bases give significantly increased binding affinity compared to either modification alone. The 2'-O-allyl modification renders oligomers incompetent as a substrate for RNase H. Thus, RNAs bound by oligomers containing 2' modifications and the invention bases can be advantageously used as probes for RNA sequences in cellular extracts containing nucleases.

20

25

Example 19

Triplex Formation Using Oligomers Containing U' and a Switchback Linker (X') Containing U'

30

The following oligomers were synthesized. All linkages were phosphodiester.

35

-110-

ON-41

5' U<sup>•</sup>C'U<sup>•</sup>C'U<sup>•</sup>U<sup>•</sup>U<sup>•</sup>U<sup>•</sup>U<sup>•</sup>C'U<sup>•</sup>U<sup>•</sup>C'U<sup>•</sup>C'U<sup>•</sup>U<sup>•</sup>U<sup>•</sup>C'X<sup>•</sup>U<sup>•</sup>U<sup>•</sup>U<sup>•</sup>U<sup>•</sup>U<sup>•</sup>U<sup>•</sup> 3'

ON-42

5' U'C'U'C'U'U'U'U'U'U'c'u'u'c'u'c'u'u'u'c'x'u'u' 3'

5 The DNA target sequence used was:

**DNA Duplex Target:** 5' AGAGAAGGGAGAAGAGAAAGAAATTTTTTTTTT 3'  
3' TCTCTTTTTTCTTCTCTTCTTTAAAAA 5'

The target sequence was introduced into an intron of the TAG gene and was used in DNase protection analysis as a linearized restriction fragment at about 2 nM. The switchback linker synthon, X', had the structure 2 shown in Figure 5-1 and was incorporated into ON-41 and ON-42 by standard H-phosphonate chemistry. The switchback linker was associated with 2 null base pairs (bases in the target that were not hybridized with). Full DNase protection of the target sequence was obtained using either ON-41 or ON-42 at a concentration less than 100 nM.

20           These results demonstrate that the invention  
bases are compatible with switchback linkers, such as the  
o-xyloso linker described here.

### Example 20

## Triple Helix Formation Using Oligomers Containing U' and Phosphorothioate Linkage

30 Triple helix formation was demonstrated using ON-5 and the phosphorothioate form of ON-5 as described in Example 16. The results show that ON-5 has a  $T_m$  of 55.8°C and the phosphorothioate derivative has a  $T_m$  of 44.9°C. Thus, the oligomer containing the phosphorothioate linkage, in combination with the bases

-111-

of the invention, is binding competent in triple helix formation.

5 The instant invention is shown and described herein in what is considered to be the most practical and preferred embodiments. It is recognized, however, that departures can be made therefrom which are within the scope of the invention, and that modifications will occur to those skilled in the art upon reading this disclosure.

10

15

20

25

30

35